
Title: APPLICATION OF S_N AND MONTE CARLO CODES

TO THE SHEBA CRITICAL ASSEMBLIFS

Author(s): R. DOUGLAS O'DELL

Submitted to: 1993 Winter Meeting of the American Nuclear Society







APPLICATION OF \mathbf{S}_N AND MONTE CARLO CODES TO THE SHEBA CRITICAL ASSEMBLIES

R. Douglas O'Dell

Nuclear Criticality Safety Group Health & Safety Division Los Mamos National Laboratory Los Alamos, NM 87545

DISCLAIMER

This report was prepared as an account of work spansored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to a specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed berein do not necessarily state or reflect those of the United States Government or any agency thereof.

Submitted for Presentation at the 1993 Winter Meeting of the American Nuclear Society San Francisco, California

APPLICATION OF \mathbf{S}_N AND MONTE CARLO CODES TO THE SHEBA CRITICAL ASSEMBLIES

R. Douglas O'Dell Los Alamos National Laboratory

The Solution High Energy Burst Assembly (SHEBA) at Los Alamos is a low enriched (1.95 wt. %) aqueous uranyl fluoride solution critical assembly. There are two SHFBA configurations, both consisting of right circular cylinders with a central control rod. The first configuration, hereafter called the old SHEBA, had a fuel solution diameter of 54.6 cm and a measured critical solution height of 36.5 cm. An improved modification, hereafter called the new SHEBA, has a fr.! solution diameter of 18.9 cm but since it is not yet operational, the critical solution height has not yet been measured. In this presentation we describe the application of the discrete ordinates (S_N) code TWODANT¹ using Hausen Roach cross sections²⁻⁴ and the MCNP Monte Carlo code¹ using continuous energy cross sections for calculating the critical solution heights for both the old and new SHEBA assemblies. We compare the code's predictions and show that a single calculation with a standard computer code may yield misleading results, especially when using a Monte Carlo code.

Initially, calculations were made on the old SHFBA with the reported critical solution height of 36.5 cm. TWODAN1 with Hansen Roach cross sections gave a k_{eff} of 0.998 and MCNP yielded a k_{eff} of 1.0020 ± 0.0022. These results appeared to be in excellent agreement with the experimental k_{eff} = ± 00. Calculations were then made on the new SHFBA with a predicted critical solution height of 11.0 cm. TWODAN1 yielded a k_{eff} of 0.9996 and MCNP gave a k_{eff} of 1.0031 ± 0.0029. Again the agreement between the two codes (and cross section data) appeared to be excellent.

We then made additional independent calculation, with MCNP in which the starting random

numbers were changed and, in some cases, the number of active cycles and number of histories per cycle were also changed. The results of these calculations are shown in the table. These results indicate that for both the old and new SHEBA, the initial MCNP $k_{
m eff}$'s appear to be on the low side of the overall uncertainty range in k_{elf} . They also show that there can be a *subconscious* bias in the person performing the calculations. In the initial runs we were expecting a $k_{
m eff}$ of unity, so we tended to mentally accept the results (within one sigma of unity) as confirming that MCNP did, in fact, agree with what we wanted the answer to be. In actuality, the MCNP results indicate a predicted $k_{
m eff}$ closer to 1.01 than to 1.00 for both SHEBAs. While these differences in $k_{
m eff}$ may not seem significant, they translate into noticable differences in critical height of the solution. For the old SHEBA, MCNP predicts a critical height of about 35.5 cm instead of 36.5 cm; for the new SHEBA, MCNP predicts a critical height of about 42 cm instead of 44 cm. The differences between the TWODANT and the MCNP predictions are, in fact, differences due to differing cross sections, that is, differences between flausen Roach 16 group data and FNDF/B V continuous energy data. The results of this work point out several items of interest. First, one must be very careful in interpreting a Monte Carlo result reported as $k_{
m eff} \pm 0.00x$. It seems to be human nature to focus on the value of $k_{
m df}$. We may recognize that there is an uncertainty band associated with that value of k_{eff} , but we usually still think in terms of the value of k_{eff} . It is suggested by the author that one develops the habit of thinking (and reporting) Monte Carlo k_{ell} as simply 2k_{ell} lies between k1 and $k2^{\circ}$ where k1 is, perhaps, $k_{\rm cab}=2\sigma$ and k2 is $k_{\rm cab}=1/2\sigma$. In actuality, this is all that we can say about the kgr from a Moute Carlo calculation. Second, it is interesting to see that with the old SIII BA, the Hansen Roach cross sections predict the actual critical height quite well while the EXDL/BA cross sections tend to underpredict the actual critical height. For the new SILLBA the predicted critical solution height free. The Hansen-Roach cross sections is similarly greater than that

predicted from the FNDF/B V cross sections used by MCNP. It will be interesting to see what the actual experimental critical height is in the new SHEBA. Finally, we note that while the k_{eff} values for a given solution height differ by less than 1 % between Hansen Roach and ENDF/B V cross sections, the difference in solution height is pronounced—about 1.2 cm.

MCNP Results for Old SHEBA (36.5 cm height) & New SHEBA (44.0 cm height)

[Old SHEBA		New SHFBA	
1	No. Active		No. Active	
Run	Histories	k _{elf} Ι· σ	Histories	k _{elf} : Εσ
<u>Initial</u>	150,000	1.0020 4. 0.0022	000,000	1.0034 ± 0.0029
\ <u>\</u>	100.000	1.0027 ± 0.0026	100.000	1.0088 ± 0.0024
1				
2	100,000	1.0062 0.0026	100,000	1.0127 0.0028
.,	20.000	Law St. Lagarit		
3	50,000	1.0072 ± 0.0031	100,000	1.0069 0.0012
1	50,000	1.0095 ± 0.0038	000,001	1.0093 0.0013
5			200,000	1.0069 1 0.0018
6			.200,000	1.0100 0.0018
l l	1		L	l. <u></u>

References

- R. E. ALCOUFFE, F. W. BRINKLEY, Jr. D. R. MARR, R. D. O'DELL, "User's Guide for TWODANT: A Code Package for Two Dimensional, Diffusion Accelerated, Neutral Particle Transport," Los Alamos National Laboratory report LA 10049 M (Rev), (February 1, 1990).
- Gordon E. HANSEN and William H. ROACH, "Six and Sixteen Group Cross Sections for Fast and Intermediate Critical Assemblies," Los Alamos Scientific Laboratory report LAMS 2543 (December 1961)
- G. I. BELL, J. J. DEVANEY, G. E. HANSEN, C. B. MILLS, W. H. ROACH, "Los Alamos Group Averaged Cross Sections," Los Alamos Scientific Laboratory report LAMS 2941 (September 1963).
- John S. HENDRICKS & Judith F. BRIESMFISTER, "The MCNP4 Monte Carlo General Purpose Radiation Transport Code," Proceedings of the International Topical Meeting on Ideances in Mathematics, Computations, and Reactor Physics, CONF 910414 (Pittsburgh, Pennsylvania, April, 1991), Vol. 5.