

CIC-14 REPORT COLLECTION
REPRODUCTION
COPY

LA-2114

~~6.3~~ C.3

LOS ALAMOS SCIENTIFIC LABORATORY
OF THE UNIVERSITY OF CALIFORNIA ○ LOS ALAMOS NEW MEXICO

SUMMARY OF NEUTRON-INDUCED FISSION CROSS SECTIONS

LOS ALAMOS NATL. LAB. LIBS



3 9338 00359 4065

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.

LA-2114
PHYSICS
(Distributed according to
TID-4500, 12th ed.)

LOS ALAMOS SCIENTIFIC LABORATORY
OF THE UNIVERSITY OF CALIFORNIA LOS ALAMOS NEW MEXICO

REPORT WRITTEN: February 25, 1957

REPORT DISTRIBUTED: May 8, 1957

SUMMARY OF NEUTRON-INDUCED FISSION CROSS SECTIONS*

by

R. L. Henkel

*This report supersedes part of LA-1714

Contract W-7405-ENG. 36 with the U. S. Atomic Energy Commission

LOS ALAMOS NATIONAL LABORATORY



3 9338 00359 4065

A recent compilation of high-energy neutron data¹ has been made for fissionable nuclei. Included were measurements of the fission cross sections for U^{233} , U^{235} , U^{238} , and Pu^{239} which were much more complete than the earlier summary given in LA-1714.² In view of the probable interest in the practical application of these measurements, it seemed desirable to distribute the results in a form which would allow the determination of cross section values with accuracies comparable with experimental uncertainties.

Most of the data are from recent measurements, which as yet are not published, and probably give the best results available at this time. The descriptions of the experiments and the uncertainties of the results are given in reference 1 and will not be repeated here. In each case, the solid line (see Figs. 1 to 4) should indicate the probable cross section values. The position of this line has been influenced by the uncertainties in the results from the various measurements.

-
1. D. W. Allen and R. L. Henkel, in Progress in Nuclear Energy, Series I, Vol. II, Pergamon Press Ltd., London (in press).
 2. H. H. Barschall and R. L. Henkel, Los Alamos Scientific Laboratory Report LA-1714, August 1954 (classified).




Fig. 1a Fission cross section of U²³³ from 0.01 to 0.1 Mev




Fig. 1b Fission cross section of U^{233} from 0.1 to 1.0 Mev




Fig. 1c Fission cross section of U^{233} from 1.0 to 10 Mev




Fig. 2a Fission cross section of U^{235} from 0.01 to 0.1 Mev




Fig. 2b Fission cross section of U^{235} from 0.1 to 1.0 Mev




Fig. 2c Fission cross section of U^{235} from 1.0 to 10 Mev




Fig. 3 Fission cross section of U^{238} from 1 to 10 Mev




Fig. 4a Fission cross section of Pu^{239} from 0.01 to 0.1 Mev




Fig. 4b Fission cross section of Pu²³⁹ from 0.1 to 1.0 Mev




Fig. 4c Fission cross section of Pu^{239} from 1.0 to 10 Mev