

CIC-14 REPORT COLLECTION  
**REPRODUCTION**  
**COPY**

LAMS-2200  
**e.3**

**LOS ALAMOS SCIENTIFIC LABORATORY**  
**OF THE UNIVERSITY OF CALIFORNIA ○ LOS ALAMOS NEW MEXICO**

---

$U^{239}$  FISSION CROSS SECTION

LOS ALAMOS NATIONAL LABORATORY



3 9338 00359 4578

## LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission to the extent that such employee or contractor prepares, handles or distributes, or provides access to, any information pursuant to his employment or contract with the Commission.

LAMS-2200  
PHYSICS AND MATHEMATICS  
(TID-4500, 13th ed., rev.)

**LOS ALAMOS SCIENTIFIC LABORATORY**  
**OF THE UNIVERSITY OF CALIFORNIA LOS ALAMOS NEW MEXICO**

REPORT WRITTEN: February 1958

REPORT DISTRIBUTED: May 16, 1958

$U^{239}$  FISSION CROSS SECTION

by

Joseph J. Devaney

Contract W-7405-ENG. 36 with the U. S. Atomic Energy Commission

All LAMS reports are informal documents, usually prepared for a special purpose and primarily prepared for use within the Laboratory rather than for general distribution. This report has not been edited, reviewed, or verified for accuracy. All LAMS reports express the views of the authors as of the time they were written and do not necessarily reflect the opinions of the Los Alamos Scientific Laboratory or the final opinion of the authors on the subject.



In the course of calculating probabilities of repeated neutron capture of  $U^{238}$  in a high density fast neutron flux, we made a crude estimate of the  $U^{239}$  fission cross section. This report gives that estimate and its basis.

We think of the process of neutron-induced fission as taking place following the capture of a neutron and the subsequent sharing of the excitation among many neutrons, from which collective modes may lead to fission itself. Ignoring all but the main features in the above concept, we recognize two gross functional forms of the fission cross section:

1. The exponential rise of the fission width as the top of the fission barrier is approached.
2. The inverse velocity dependence of the average compound nucleus cross section.

We therefore look for a crude cross section dependence below the fission barrier of the form

$$\sigma_F \propto \sigma_{FCN} \cdot \Gamma_F$$

where  $\sigma_F$  is the fission cross section,  $\sigma_{FCN}$  is the cross section for formation of the compound nucleus,  $E_n$  is the neutron energy, and  $\Gamma_F$  is the fission width. We shall expect  $\sigma_{FCN} \propto E_n^{-1/2}$  and  $\Gamma_F$  to be an exponential.

The following considerations were, therefore, incorporated in Fig. 1:

- A. The well-known fact that for given element, both even mass numbered isotopes and the odd mass numbered isotopes separately show a

decrease in "threshold" energy with increasing mass number.<sup>1-4</sup> We therefore estimate that  $U^{239}$  has a sharp positive rise in the fission cross section at about 0.47 Mev.

B. The empirical variation of the "plateau" value of the fission cross section with  $Z^{4/3}/A$  as put forward by Barschall and Henkel.<sup>5</sup> Our estimate from this work is that  $U^{239}$  has a "plateau" value of about 0.25 barn.

C. The simplest exponential law for  $\Gamma_F$  is  $e^{a(E-E_0)}$ . We estimate a by comparison with an extrapolation from the thresholds<sup>6,7</sup> of  $U^{234}$ ,  $U^{236}$ ,  $U^{238}$ ,  $Np^{237}$ , and  $Am^{241}$ , using the more complete form of D below to arrive at  $a = 8.3 \text{ Mev}^{-1}$ . Compare Seaborg.<sup>3</sup>

D. Putting in the thermal average value<sup>7</sup> of 15 barns at  $0.0253 \times 10^{-6}$  Mev and ignoring resonance structure, we estimate  $E_0$  from the formula

$$\langle \sigma_F \rangle = 0.25 \sqrt{\frac{E_0}{E}} e^{8.3(E-E_0)}$$

- 
1. Philip Morrison in Experimental Nuclear Physics, E. Segrè, editor, Vol. II, page 136, John Wiley & Sons, Inc., New York (1953). Not used quantitatively. Values too high for  $U^{235}$ .
  2. Neutron binding energies, etc., were taken from A. G. W. Cameron, Atomic Energy of Canada Limited, Chalk River Project Report CRP-690 (March, 1957).
  3. G. T. Seaborg, Phys. Rev. **88**, 1429 (1952).
  4. Alex E. S. Green, Nuclear Physics, International Series of Physics, McGraw-Hill Book Company, Inc., New York (1955).
  5. H. H. Barschall as reported in R. L. Henkel, Los Alamos Scientific Laboratory Report LA-2122 (June, 1957).
  6. R. L. Henkel, Los Alamos Scientific Laboratory Report LA-2122 (June, 1957), and private communications (1956-57).
  7. Donald J. Hughes and Robert B. Schwartz, "Neutron Cross Sections," Brookhaven National Laboratory Report BNL-325, Supplement I, (January, 1957).

and we find  $E_0 = 0.521$  Mev. This result is in good agreement with the independent estimate of A which gives a "threshold" of the order of 0.47 Mev. Figure 1 shows that the above formula is in fortuitously good agreement.

E. The graph, therefore, incorporates the formula

$$\langle \sigma_F \rangle \text{ (barns)} = 0.25 [0.521/E(\text{Mev})]^{1/2} \exp[8.3 (E - 0.521)]$$

below  $E = 0.38$  Mev with a plateau of 0.25 barn at about 2 to 4 Mev, the two being joined by a smooth curve of no physical significance whatever.




Fig. 1  $\sigma_F$  of  $U^{239}$  vs neutron energy.