

# **LEGIBILITY NOTICE**

**A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.**

**Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.**

LA-UR -89-2541

CONF-890928--1

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA-UR--89-2541

DE89 015288

TITLE FORMATION, CHARACTERIZATION, AND STABILITY OF PLUTONIUM(IV)

COLLOID: A PROGRESS REPORT

AUTHOR(S) David E. Hobart, David E. Morris, Phillip D. Palmer and  
Thomas W. Newton

Nuclear and Radiochemistry Group (INC-11)  
Isotope and Nuclear Chemistry Division  
Los Alamos National Laboratory

SUBMITTED TO

American Nuclear Society Topical

Proceedings of "FOCUS 89: Nuclear Waste Isolation in the  
Unsaturated Zone," Las Vegas, Nevada, September 18-21, 1989

#### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

 Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

11/9

[For invited publication in the American Nuclear Society Topical: *Proceedings of "FOCUS 89: Nuclear Waste Isolation in the Unsaturated Zone,"* Las Vegas, Nevada, September 18-21, 1989.]

LA-UR-89-XXXX

## **FORMATION, CHARACTERIZATION, AND STABILITY OF PLUTONIUM(IV) COLLOID: A PROGRESS REPORT\***

D. E. Hobart, D. E. Morris, P. D. Palmer,  
and T. W. Newton

Isotope and Nuclear Chemistry Division  
Los Alamos National Laboratory  
Los Alamos, New Mexico, 87545 USA

*This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees makes any warranty, express or implied or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer or otherwise does not necessarily constitute or imply its endorsement, recommendation or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.*

(WBS 1.2.3.4.1.4.A/QA Level 1)

\*This work was supported by the Yucca Mountain Project Office as part of the Civilian Radioactive Waste Management Program. This project is managed by the U.S. Department of Energy, Nevada Operations Office. Support was also provided by the Division of Chemical Sciences, Office of Basic Energy Sciences, U.S. Department of Energy under contract W-7405-ENG-36 with The University of California

## FORMATION, CHARACTERIZATION, AND STABILITY OF PLUTONIUM(IV) COLLOID: A PROGRESS REPORT

D. E. HOBART, D. E. MORRIS, P. D. PALMER, and T. W. NEWTON\*  
Isotope and Nuclear Chemistry Division  
Los Alamos National Laboratory  
Los Alamos, New Mexico 87545 USA  
(505) 667-9313

### ABSTRACT

Plutonium is expected to be a major component of the waste element package in any high-level nuclear waste repository. Plutonium(IV) is known to form colloids under chemical conditions similar to those found in typical groundwaters. In the event of a breach of a repository, these colloids represent a source of radionuclide transport to the far-field environment, in parallel with the transport of dissolved waste element species. In addition, the colloids may decompose or disaggregate into soluble ionic species. Thus, colloids represent an additional term in determining waste element solubility limits. A thorough characterization of the physical and chemical properties of these colloids under relevant conditions is essential to assess the concentration limits and transport mechanisms for the waste elements at the proposed Yucca Mountain Repository site. This report is concerned primarily with recent results obtained by the Yucca Mountain Project (YMP) Solubility Determination Task (WBS 1.2.3.4.1.4.A) pertaining to the characterization of the structural and chemical properties of Pu(IV) colloid. Important results will be presented which provides further evidence that colloidal plutonium(IV) is structurally similar to plutonium dioxide and that colloidal plutonium(IV) is electrochemically reactive.

### INTRODUCTION

It has long been known that aqueous Pu(IV) forms colloids under appropriate chemical conditions.<sup>1</sup> This material is often called "polymeric" Pu(IV) but colloid or colloidal is the preferable label so as not to imply necessarily that the material has a chain-like structure. The colloid can form solution-like *cols* that are optically clear, display a characteristic absorption spectrum, and do not settle out on long standing.<sup>1-3</sup>

\* Guest scientist of the Isotope and Nuclear Chemistry Division

In noncomplexing acid solutions with concentrations greater than about 0.05 M, Pu(IV) is predominantly aquated  $\text{Pu}^{4+}$ . This highly charged ion hydrolyzes readily. In 0.05 M acid and unit ionic strength,  $\text{Pu}(\text{OH})^{3+}$  and  $\text{Pu}^{3+}$  have about equal concentrations. At lower acid concentrations further hydrolysis is expected, giving the ions  $\text{Pu}(\text{OH})_2^{2+}$ ,  $\text{Pu}(\text{OH})_3^+$  and neutral species such as  $\text{Pu}(\text{OH})_4^0$ . Formation constants have been estimated for these species,<sup>4</sup> but measurements have been made only for the  $\text{Pu}(\text{OH})^{3+}$  species.<sup>5</sup> An important reason for this lack of data is that colloidal Pu(IV) forms quite rapidly in the pH region where the more highly hydrolyzed species can be studied. In addition, aqueous Pu(IV) is unstable with respect to Pu(III) and Pu(V) as shown, for example, in the oxidation state stability field diagram published by Allard et al.<sup>6</sup>

Pu(IV) colloid is of particular interest in nuclear waste management because it is one of the dominant forms of plutonium under chemical conditions similar to those found in the environment. There is also an interest in this material with regard to its formation in process streams and in its utilization in the production of nuclear fuel elements by *sol-gel* processes.<sup>7</sup> We are currently involved in an ongoing program to investigate the formation and stability of Pu(IV) colloid and to characterize the chemical and physical properties of the colloid. In this report, some recent results are discussed pertaining to the chemical and structural characterization of Pu(IV) colloid using electronic absorption and diffuse reflectance spectroscopies and electrochemical and redox titrimetric methods.

One of the most important, yet relatively unexplored, aspects in the chemical characterization of Pu(IV) colloid is the determination of the structure of the colloid. Several direct structural probes have been utilized. However, in the published structural studies,<sup>2,3</sup> it has been necessary for experimental reasons to examine the Pu(IV) colloid in a dried form.

Recently, attempts have been made in our laboratories to deduce structural information for  $\text{Pu(IV)-colloid}$  suspended in solution utilizing electronic spectroscopy as a probe by comparing spectral data for the colloid with that obtained for other  $\text{Pu(IV)}$  complexes.

Information on the redox reactivity of  $\text{Pu(IV)}$  colloid, i.e., the potentials at which oxidation and reduction occur and the rates and mechanisms of these processes, is of fundamental importance to the characterization of this species because redox reactions may represent viable mechanisms of colloid degradation to give dissolved ionic species with enhanced environmental mobility. Further, results relating to redox reactivity will be of great value in defining the general reactivity patterns for the colloid so that other pertinent chemical kinetic phenomena can be interpreted more readily. Finally, it may be possible to utilize voltammetric redox techniques to obtain results relating to the physical properties of the colloid. Such data would be useful to corroborate the results obtained using other techniques.

## EXPERIMENTAL

Colloidal sols of  $\text{Pu(IV)}$  and the solid complexes  $\text{Pu}_2\text{C}_2\text{O}_7 \cdot 6\text{H}_2\text{O}$  and  $\text{Pu}_2\text{SO}_4 \cdot 4\text{H}_2\text{O}$  were prepared and purified according to standard procedures described in detail elsewhere.<sup>8</sup> Samples of high-fired  $\text{PuO}_2$  were obtained from the Materials and Science Technology Division of Los Alamos National Laboratory. The  $\text{PuF}_4$  sample was generously provided by J. D. Farr of Los Alamos National Laboratory. The electronic spectroscopic studies were carried out using a Varian Cary Model 17D UV-Vis-NIR spectrophotometer. For the colloidal sols, the instrument was operated in the standard absorption mode. However, because of the insolubility of  $\text{PuO}_2$  and  $\text{Pu}_2\text{C}_2\text{O}_7$  and the opacity of these materials as finely divided solids it was necessary to obtain the electronic spectra for these samples using diffuse reflectance methods. The electronic spectra of  $\text{PuF}_4$  and  $\text{Pu}_2\text{SO}_4$  were also obtained by diffuse reflectance. The custom-designed diffuse reflectance attachment for the Varian Cary spectrophotometer was obtained on loan from Dr. Jack Young of the Analytical Chemistry Division of Oak Ridge National Laboratory. With the reflectance apparatus in place, the instrument was operated in the usual fine-beam absorption mode. Neutral density screen filters were placed in the reference beam of the spectrophotometer and the absorbance was adjusted electronically as necessary to keep the signal output within the absorbance range of the instrument.

The voltammetric investigations of  $\text{Pu(IV)}$  colloid were conducted in a microvolume electrolysis cell described in detail elsewhere.<sup>9</sup> The bulk electrolytic

studies were done in either a long path-length spectroelectrochemical cell or a standard three-electrode electrolysis cell. Working electrodes were fabricated from platinum for the oxidation studies and mercury for the reduction studies. These experiments were controlled with an EG&G Princeton Applied Research Corporation Model 273 Potentiostat interfaced to an IBM PC/AT using the HEADSTART software package or with a Model 173 Potentiostat and Model 175 Universal Programmer. Tripartite investigations were monitored spectrophotometrically using the Cary Model 17D in the standard absorption mode.

## RESULTS AND DISCUSSION

### A Spectroscopic Characterization

When the electronic absorption spectrum of  $\text{Pu(IV)}$  colloid is compared with that obtained for the aquated  $\text{Pu}^{4+}$  ion (Figure 1), it is clear that there are numerous differences between these spectra.




FIGURE 1. ABSORPTION SPECTRA OF AQUEOUS  $\text{Pu(IV)}$  AND  $\text{Pu(IV)}$  COLLOIDAL SOL.

This suggests that there are significant differences in the environments of these two tetravalent plutonium species. In particular, the most prominent visible absorption bands for  $\text{Pu(IV)}$  colloid occur at  $\sim 750$ ,  $\sim 625$ , and  $\sim 740$  nm, whereas the major visible absorption bands for aquated  $\text{Pu}^{4+}$  are at  $\sim 470$ ,  $\sim 660$ , and  $\sim 800$  nm. In contrast to these major differences between the spectra for the colloid and aquated  $\text{Pu}^{4+}$ , the band maxima for  $\text{Pu(IV)}$  colloid are strongly correlated with the band maxima in the diffuse reflectance spectrum of high-fired  $\text{PuO}_2$  in the visible and near-IR spectral regions (Figure 2).




FIGURE 2 ABSORPTION SPECTRUM OF  $\text{Pu(IV)}$  COLLOIDAL SOL AND DIFFUSE REFLECTANCE SPECTRUM OF HIGH FIRED PLUTONIUM DIOXIDE

It should be emphasized that diffuse reflectance spectroscopy provides the same information as absorption spectroscopy with respect to the energies of the transitions between energy levels in the analyte. Thus, the locations of the bands in the UV-visible near infrared spectral region arising from electronic transitions are expected to be the same for any given sample whether determined by absorption spectroscopy or diffuse reflectance spectroscopy. However, the mechanisms that determine the intensities in these spectral bands are quite different for diffuse reflectance than for absorption. Therefore, there will be no discussion of intensities in this report.

The number of bands in the electronic spectrum of an actinide complex and the positions of the band maxima depend to a large extent on the environment of the actinide ion. For this reason, electronic spectroscopy can be used as a sensitive structural probe. Tetravalent plutonium has a ground state electronic configuration (Radon core)  $5f^4$  and the electronic spectrum of this ion is dominated by  $f-f$  transitions. In the free ion case

(i.e. no complexation of  $\text{Pu(IV)}$ ); these transitions are strictly parity forbidden. Similarly, for complexed  $\text{Pu(IV)}$  in a sufficiently high symmetry environment, the electronic transitions from the ground vibrational level of the ground electronic state to the ground vibrational level of the excited electronic state (the so called electronic origin transitions) will also be forbidden. In this case the observed spectral bands (neglecting any crystal field splittings) are due to transitions to other excited vibrational levels in the excited electronic state or from other excited vibrational levels of the ground electronic state. Such transitions are referred to as vibronic transitions. For  $\text{Pu(IV)}$  in a low symmetry environment, the electronic spectrum can contain both electronic origin transition and vibronic transitions.

A thorough interpretation of the electronic spectra of  $\text{Pu(IV)}$  complexes would require a complete group theoretical treatment and additional experimental data. Such a treatment is beyond the scope of the present work. However, the implication of the above discussion is that significant differences in the electronic spectra of  $\text{Pu(IV)}$  complexes are

expected if the complexes have different site symmetry for the Pu(IV) ion and/or if they have different vibrational energy level spacings, as would result from different complexing ligands or differing bond strengths for bonds to the same ligand.

The diffuse reflectance spectra of the other Pu(IV) complexes, which possess various symmetries and differing vibrational energies, were obtained to explore further this correlation between the electronic spectra and the structure in Pu(IV) complexes. The crystallographic symmetry of the oxide complex is face-centered cubic, that of the oxalate complex is triclinic and that of the sulfate complex is orthorhombic. The symmetry of the fluoride complex depends on the hydration number and is unknown for the sample employed in this study. However, it is most likely the anhydrous salt which is monoclinic. It should be emphasized that the site symmetry of the Pu(IV) ion in these complexes is not necessarily the same as the crystallographic symmetry, but the crystallographic symmetry is a useful initial point for comparison. The spectra of these complexes are shown in Figures 3 and 4, with the spectrum of PuO<sub>2</sub> serving as a reference for comparison. These spectral data clearly reveal the variations in the electronic spectra that result from differences in the symmetry environment and vibrational energies.

The excellent correlation between the spectrum of Pu(IV) colloid and that of high fired plutonium




FIGURE 3 DIFFUSE REFLECTANCE SPECTRA OF PLUTONIUM DIOXIDE, Pu(IV) FLUORIDE, AND Pu(IV) SULFATE.




FIGURE 4 DIFFUSE REFLECTANCE SPECTRA OF PLUTONIUM DIOXIDE AND Pu(IV) OXALATE.

dioxide (Fig. 2) can be compared to the variations between the spectra shown in Figures 3 and 4. This large degree of coincidence between the colloid spectrum and the oxide spectrum is strong evidence that the structure of the colloid is very similar to that of high fired PuO<sub>2</sub>. This conclusion has been suggested previously by others on the basis of x-ray<sup>1</sup> and electron<sup>2</sup> diffraction studies. Furthermore, because high fired PuO<sub>2</sub> is a high symmetry complex, the observed electronic bands are predominantly due to vibronic transitions. This implies that conclusions regarding the structural similarity of the colloid and PuO<sub>2</sub> can be extended to include similarities in the vibrational energies in addition to similarities in the site symmetry of the Pu(IV) ion. This conclusion is also supported by previous infrared spectral studies wherein it was shown that the two prominent vibrational modes of PuO<sub>2</sub> are also seen at the same energy in the colloid.<sup>10</sup> Thus, these electronic spectral results indicate that Pu(IV) colloid, when suspended in solutions, is structurally quite similar to solid high fired PuO<sub>2</sub>.

A photoacoustic spectrum of PuO<sub>2</sub> has been published by Heinrich et al.<sup>11</sup> however, it lacked sufficient resolution and detail to allow observation of its resemblance to the spectrum of Pu(IV) colloid.

## B Electrochemical Characterization

The electrochemical behavior of dissolved plutonium in its four readily accessible oxidation states is well characterized<sup>12</sup> and can be interpreted within the framework of existing simple theories of solution phase thermodynamics and kinetics. In contrast, there are very few reports concerning the application of electrochemical methods to the study of Pu(IV) colloid. Presumably, this paucity of data can be attributed in large part to the inherent complexity in investigating redox reactivity in distinctly heterogeneous media such as colloidal suspensions. Despite the anticipated complexity in the redox reactions of Pu(IV) colloid, the dominant electrochemical processes are still expected to be reduction to a dissolved Pu(III) species and oxidation to a dissolved Pu(VI)O<sub>2</sub><sup>2+</sup> species. This latter process may also involve a Pu(V) species as an intermediate.

The initial experiments were designed to test these hypotheses. To determine the product of the reduction of Pu(IV) colloid, Zn-Hg amalgam, with a potential of 0.76 V versus the normal hydrogen electrode (NHE), was chosen as a reducing agent. Using a specially designed spectroelectrochemical cell,<sup>13</sup> a solution of Pu(IV) colloidal suspension in dilute perchloric acid was stirred in the presence of excess Zn(Hg). The course of the reaction was monitored spectrophotometrically by detecting the diminution of the spectrum of Pu(IV) colloid and the growth of the spectrum of the reduction product. The product spectrum was readily identified as that of dissolved aquated Pu<sup>3+</sup>. This reduction reaction was determined to follow simple first-order kinetics with a half time of ~40 min. In a related experiment, the Zn(Hg) amalgam was replaced with a potentiostatically controlled mercury cathode and the rate of reduction of the colloid was monitored spectrophotometrically as a function of applied potential. In this case, it was determined that a significant reduction rate occurs even at ~0.46 V versus NHE. The reaction appears to reach a maximum rate at ~0.76 V versus NHE and this rate is maintained at more negative potentials. To determine the product of the oxidation of Pu(IV) colloid, the reaction of a stirred solution of colloid in contact with a potentiostatically controlled platinum gauze anode was studied. Here, too, the progress of the reaction was followed spectrophotometrically. In this case, it was found that no significant rate of colloid oxidation takes place until the potential reaches ~+1.6 V versus NHE. The product of the oxidation was identified as Pu(VI)O<sub>2</sub><sup>2+</sup>. No Pu(V)O<sub>2</sub><sup>+</sup> was detected, but the existence of this species as a short-lived intermediate in this reaction cannot be excluded because the time scale of the assay was rather long.

Investigations of the oxidation of Pu(IV) colloid by cerium(IV) in perchloric acid were initiated several years ago in our laboratories. The reaction was found to be quite complicated and was not pursued. A recent report of similar studies in nitric acid<sup>14</sup> led to a reinvestigation of these studies. Experiments have now been conducted in both nitric and perchloric acids for several colloid preparations which differ in particle size. The potential of the Ce(IV)-III couple is ~+1.6 V in nitric acid and ~+1.7 V in perchloric acid versus NHE. The course of the oxidation reaction was followed by measuring the spectrophotometric absorbance of the product Pu(VI)O<sub>2</sub><sup>2+</sup> peak near 830 nm. The results indicate that the initial rate of colloid oxidation is quite fast in all cases with the rate being faster in nitric acid than in perchloric acid. However, the reaction goes to completion only in nitric acid. In perchloric acid, the reaction becomes "stalled" after between 30 and 70% of the total Pu(IV) has reacted. This variation in the extent of completion appears to depend on particle size, acid concentration, and Ce(IV) concentration in a complicated way. No obvious trends in these reactivity patterns have been determined as yet. Representative kinetic data for these experiments, in the form of a Powell plot, are shown in Figure 5. The open circles are experimental data for a Pu(IV) colloid sample in dilute nitric acid with a most probable particle diameter of 2.1 nm. The filled circles are data for a Pu(IV) colloid sample in dilute perchloric acid with a most probable particle diameter of 25.5 nm. The dashed line is the theoretical curve for a first-order reaction and the solid line is the theoretical curve for a second-order




FIGURE 5. POWELL PLOTS FOR THE OXIDATION REACTION OF Ce(IV) WITH Pu(IV) COLLOID



reaction. Note that the data in nitric acid appear to follow theoretical second-order kinetic behavior. This is a very surprising result because it suggests that the rate law is second-order in  $\text{Pu(IV)}$  colloid concentration. A simple mechanism consistent with this finding cannot be formulated. The data from experiments in perchloric acid do not appear to follow any simple rate law. These results are dramatically different from those reported previously<sup>12</sup> in which a first-order dependence on colloid concentration was found in nitric acid. The present results also indicate that the small particle-size colloid samples react more rapidly than the large particle-size samples in nitric acid. These studies are continuing in hopes of unraveling these puzzling kinetic phenomena.

The voltammetric studies have used two standard techniques: cyclic voltammetry and chronoamperometry. Both techniques provide a much shorter time scale interrogation of the redox reactions than the methods discussed above. All of these experiments have been done in dilute hydrochloric acid and in the absence of stirring so that diffusion and migration are the only means of delivery of the colloid to the electrode. No oxidative voltammetric activity was detected for  $\text{Pu(IV)}$  colloid by either technique for applied potentials as positive as +1.6 V versus NHE. This suggests that the rate of colloid oxidation, even at potentials approaching values at which water is oxidized, is quite slow. Reductive voltammetric activity is observed for the colloid by both techniques in the potential range from  $\sim -0.9$  to  $-1.2$  V versus NHE. This voltammetric behavior is distinctly different from the usual behavior exhibited by dissolved species. A series of chronoamperograms as a function of potential is shown in Figure 6. This sample has a most probable particle diameter of 2.1  $\mu\text{m}$ . The potentials for curves 1-4 are  $-1.20$ ,  $-1.25$ ,  $-1.30$ , and  $-1.35$  V versus a  $\text{Ag/AgCl}$  reference electrode respectively. Note that these curves exhibit a current peak at times greater than  $t=0$  instead of the usual  $t^{-1/2}$  decay for  $t>0$ . This novel behavior may be attributable to the diffusion of a partially reduced colloid particle back to the electrode surface for further reduction.

The chronoamperometric data have been used to construct the current versus potential curves shown in Figure 7. Curve 1 is for a colloid sample with a most probable particle diameter of 2.1  $\mu\text{m}$  and curve 2 is for a sample with a most probable particle diameter of 25.5  $\mu\text{m}$ . These curves show that the rate of  $\text{Pu(IV)}$  colloid reduction increases as the potential is made more negative as expected. However, the characteristic plateau in these curves at the more negative potentials is not observed. This suggests that the electron transfer step itself is kinetically quite slow.




FIGURE 6 CHRONOAMPEROGRAMS FOR THE REDUCTION OF  $\text{Pu(IV)}$  COLLOID IN DILUTE HYDROCHLORIC ACID AT A MERCURY CATHODE




FIGURE 7 CURRENT-VOLTAGE CURVES CONSTRUCTED FROM CHRONOAMPEROMETRIC DATA FOR THE REDUCTION OF  $\text{Pu(IV)}$  COLLOID IN DILUTE HYDROCHLORIC ACID AT A MERCURY ELECTRODE

Further, at any given potential, the rate of reduction of small particle-size colloid samples is much greater than that of large particle-size samples. This behavior can be attributed in part to the greater diffusivity of the smaller particles.

## CONCLUSIONS

Additional structural investigations of Pu(IV) colloid, both suspended in solution and isolated from solution, are planned to support the electronic spectral data presented here. X-ray diffraction and absorption methods and Raman spectroscopy will be exploited for these studies. The x-ray diffraction and Raman spectroscopic probes will be used to provide specific structural information. The Raman technique will be used to probe the lattice vibrational modes of the colloid in order to gain information on crystallinity and long range order in the colloid. In addition, the Raman spectral results will be used to obtain molecular structural information (e.g., the strength of oxygen bonding to plutonium) for the colloid through examination of the internal vibrational modes. To fully understand both x-ray diffraction and Raman spectral results, it will be necessary to examine certain other Pu(IV) model compounds as was done with the electronic spectral results reported here.

The preliminary results pertaining to the redox reactivity of Pu(IV) colloid reveal the anticipated complexity in the reactions of this important species. The potentials at which the colloid is oxidized and reduced are well removed from those for dissolved Pu(IV) and reflect the stability of the colloidal form of Pu(IV). Although these potentials are outside the range expected under normal environmental conditions, oxidation state stabilities of dissolved plutonium species are strongly influenced by coordination to environmentally ubiquitous complexants (such as carbonate ions), and similar effects may exist for colloidal Pu(IV). The unusual kinetic results for the Ce(IV) oxidation reaction and the novel voltammetric behavior for the colloid reduction suggest that particle size and solution composition play strong roles in determining reactivity. The results presented here have merely scratched the surface of this challenging problem and much additional work will be required to gain the necessary understanding of this system.

## ACKNOWLEDGEMENTS

This work was supported by the Yucca Mountain Project Office as part of the Civilian Radioactive Waste Management Program. This project is managed by the U.S. Department of Energy, Nevada Operations Office. Support was also provided by the Division of Chemical Sciences, Office of Basic Energy Sciences, U.S. Department of Energy under contract W-7405-ENG-36 with The University of California.

## REFERENCES

1. B. B. CUNNINGHAM, in *The Transuranic Elements*, J. J. Katz and G. T. Seaborg, eds., National Nuclear Energy Series (McGraw Hill, New York, 1954) Div. IV, Vol. 14A, Chapter 10, p. 371.
2. M. H. LLOYD and R. G. HAIRE, "The Chemistry of Plutonium in Sol-Gel Processes," *Radiochimica Acta* **25**, 139 (1978).
3. D. W. OCKENDEN and G. A. WELCH, "The Preparation and Properties of Some Plutonium Compounds. Part V," *Journal of the Chemical Society* **3358** (1956).
4. C. F. BAES and R. E. MESMER, *The Hydrolysis of Cations*, John Wiley and Sons, New York, 1976, p. 190.
5. S. W. RABIDEAU, "The Hydrolysis of Plutonium(IV)," *Journal of the American Chemical Society*, **79**, 3675 (1957).
6. B. ALLARD, H. KIPATSI, and J. O. LILJENZIN, "Expected Species of Uranium, Neptunium, and Plutonium in Neutral Aqueous Solutions," *Journal of Inorganic and Nuclear Chemistry*, **42**, 1015 (1980).
7. A. H. ZELTMANN, Los Alamos National Laboratory, private communication of unpublished results, 1973.
8. D. E. HOBART, D. E. MORRIS, and P. D. PALMER, "Formation, Characterization, and Stability of Plutonium(IV)-Colloid," Los Alamos National Laboratory report LA-UR-87-3505 (1987).
9. D. E. HOBART, D. E. MORRIS, P. D. PALMER, R. G. HAIRE, and J. R. PETERSON, "Spectroscopic and Redox Properties of Berkelium in Complexing Aqueous Carbonate and Citrate Solutions," *Radiochimica Acta*, in press.
10. L. M. TOTH, and H. A. FRIEDMAN, "The IR Spectrum of Pu(IV) Polymer," *Journal of Inorganic and Nuclear Chemistry* **40**, 807 (1978).
11. G. HEINRICH, H. GÜSTEN, and H. J. ACHE, "Photoacoustic Spectroscopy of Actinide Oxides," *Journal of Less-Common Metals*, **122**, 25 (1986).
12. S. W. RABIDEAU, "Thermodynamic Functions and Formal Potentials of the Plutonium(V)-(IV) Couple," *Journal of the American Chemical Society* **78**, 2705 (1956).
13. D. J. SAVAGE and T. W. KYFFIN, "Oxidative Breakdown of Polymeric Plutonium(IV) Hydroxide by Cerium(IV)," *Polyhedron* **5**, 743 (1986).