

**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE
(DEPARTMENT OF PURE MATHEMATICS)

ZW 195/83

JUNI

J. VINÁREK

PROJECTIVE MONADS AND EXTENSIONS OF FUNCTORS

kruislaan 413 1098 SJ amsterdam

Printed at the Mathematical Centre, Kruislaan 413, Amsterdam, The Netherlands.

The Mathematical Centre, founded 11 February 1946, is a non-profit institution for the promotion of pure and applied mathematics and computer science. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

1980 Mathematics subject classification: Primary 18C20, Secondary 18B20

Copyright © 1983, Mathematisch Centrum, Amsterdam

Projective monads and extensions of functors

by

Jirí Vinárek ^v*)

ABSTRACT

We present a class of monads with the property that any functor has an extension to the corresponding Kleisli category. These topics are connected with fuzzy-theories used for a categorical description of fuzzy-automata.

KEY WORDS & PHRASES: *set-functor, hom-functor, projective monad, Kleisli category, distributive laws, free semigroup, free monoid*

*) The author was supported by a grant of the Netherlands ministry of Education and Science during his research stay at the Mathematical Centre in Amsterdam.

Adress: Matematicko-fyzikální fakulta, Karlovy University,
Sokolovská 83, 186 00 Praha 8, Czechoslovakia.

0. INTRODUCTION

Category theory can be used to describe efficiently constructions in automata theory. It has led to a great development of studying automata using categorical methods (see e.g. [Ad], [AdTr], [ArM], [BuH], [G], [Th], [Tr1], [Tr2], etc.). Besides deterministic automata, also nondeterministic and fuzzy-automata are studied.

Whereas a deterministic automaton has a state-transition function $Q \times I \rightarrow Q$ (Q is a set of states, I - a set of inputs) a nondeterministic one has a state-transition function $Q \times I \rightarrow PQ$ (PQ is the power-set of states) assigning to each state a set of possible successors. Of course, there is a canonical embedding of a set Q into its power-set PQ and there is also a natural transformation $P^2 \rightarrow P$. This situation was generalized by the definition of monad and its Kleisli category which is entirely coextensive with fuzzy-theories as is proved in [M].

In [ArM], M.A. Arbib and E.G. Manes studied a problem when a functor $F: \underline{C} \rightarrow \underline{C}$ could be extended to the Kleisli category of a given monad. They found a sufficient and necessary condition for existence of such an extension. Their condition is analogous to the Beck distributive laws between monads (see [Be]). Therefore, the term "distributive laws" is used for these diagrams as well. It is, however, sometimes quite difficult to decide whether there exist distributive laws for a given monad and a given functor, even for some very natural monads. Such a very natural monad is the monad corresponding to the variety of monoids (i.e. semigroups with units). As was proved in [V1], this monad does not satisfy distributive laws with respect to the very "simple" functor $\text{Hom}(2, -)$ assigning to each set X its square $X \times X$. M.A. Arbib and E.G. Manes proved in [ArM] that set-functors $- \times X$ have extensions to the Kleisli category of any monad. On the other hand, we present a class of monads (called projective monads) with the property that any functor has an extension to the corresponding Kleisli category. This general result is applied to the category of free semigroups. We also present a survey of results connected with extensions of functors to the category of free monoids.

1. PRELIMINARIES

First, recall some definitions and notations:

1.1. Let \underline{C} be a category, $T: \underline{C} \rightarrow \underline{C}$ a functor, $\text{Id}: \underline{C} \rightarrow \underline{C}$ the identity functor, $\eta: \text{Id} \rightarrow T$, $\mu: T^2 \rightarrow T$ natural transformations. $\underline{T} = (T, \eta, \mu)$ is called a *monad* iff the following diagrams commute:

$$\begin{array}{ccc}
 T^3 & \xrightarrow{T\mu} & T^2 \\
 \mu T \downarrow & & \downarrow \mu \\
 T^2 & \xrightarrow{\mu} & T
 \end{array}
 \qquad
 \begin{array}{ccccc}
 & & T^2 & & \\
 \eta T \nearrow & & \downarrow \mu & & T\eta \searrow \\
 \text{Id} T & \cong & T & \cong & T \text{Id}
 \end{array}$$

1.2. The *Kleisli category* $\underline{C}(\underline{T})$ is defined as follows: the class of its objects is $\text{obj } \underline{C}(\underline{T}) = \{X(\underline{T}) \mid X \in \text{obj } \underline{C}\}$; $f(\underline{T}): X(\underline{T}) \rightarrow Y(\underline{T})$ is a morphism in $\underline{C}(\underline{T})$ if $f: X \rightarrow TY$ is a morphism of \underline{C} . Given $f(\underline{T}): X(\underline{T}) \rightarrow Y(\underline{T})$, $g(\underline{T}): Y(\underline{T}) \rightarrow Z(\underline{T})$, the corresponding composition is $g(\underline{T}) * f(\underline{T}) = (\mu_Z \circ Tg \circ f)(\underline{T})$.

1.3. a) Denote $\text{Sem} = (S, i, s)$ the monad which assigns to each set A the semigroup of words created by A , i.e. the free semigroup over A .

$(SA = \{a_1 a_2 \dots a_n \mid n \in \{1, 2, \dots\}, a_i \in A \text{ for } i = 1, 2, \dots, n\}, i_A(a) = a, s_A((a_{11} \dots a_{1k(1)}) \dots (a_{n1} \dots a_{nk(n)})) = a_{11} \dots a_{1k(1)} a_{21} \dots a_{n1} \dots a_{nk(n)}$ whenever $k(1), \dots, k(n) \geq 1$.)

b) Denote $\text{Mon} = (M, e, m)$ the monad which assigns to each set the monoid of words created by A , i.e. the free monoid over A . ($MA = SA \cup \{\Lambda\}$ where Λ is the empty word, $e_A(a) = a, m_A((a_{11} \dots a_{1k(1)}) \dots (a_{n1} \dots a_{nk(n)})) = a_{11} \dots a_{1k(1)} a_{21} \dots a_{n1} \dots a_{nk(n)}$ whenever $k(1), \dots, k(n) \geq 0$.)

1.4. DEFINITION (Arbib-Manes, [ArM]). Let \underline{C} be a category, $F: \underline{C} \rightarrow \underline{C}$ a functor, (T, η, μ) a monad, F is said to satisfy *distributive laws* over (T, η, μ) if to each object A of \underline{C} a morphism $\lambda_A: FTA \rightarrow TFA$ is assigned such that the following two diagrams commute for any A and $\alpha: A \rightarrow TB$:

(1)

$$\begin{array}{ccc}
 \text{FTA} & \xrightarrow{\lambda_A} & \text{TFA} \\
 \uparrow F\eta_A & & \uparrow \eta_{FA} \\
 & \text{FA} &
 \end{array}$$

(the first distributive law)

(2)

$$\begin{array}{ccc}
 \text{FTA} & \xrightarrow{\lambda_A} & \text{TFA} \\
 \downarrow F\alpha^+ & & \downarrow (\lambda_B \circ F\alpha)^+ \\
 \text{FTB} & \xrightarrow{\lambda_B} & \text{TFB}
 \end{array}$$

where $\alpha^+ = \mu_B \circ T\alpha$.

1.5. REMARK. A functor F can be extended to the Kleisli category $\underline{C}((T, \eta, \mu))$ iff it satisfies distributive laws.

PROOF. is given in [ArM].

1.6. a) $\text{Hom}(A, -)$ denotes a functor which assigns to each set X the set of functors $\{f: A \rightarrow X\}$.

b) A *symmetric hom-functor* $\text{Sym}(A, -)$ is a factorfunctor $\text{Hom}(A, -)/\sim$ where $f \sim g$ whenever there exists a bijection $b: A \rightarrow A$ such that $fb = g$.

1.7. REMARK. Symmetric hom-functors are a special case of tree-group varieties, investigated by V. Trnková and J. Adámek (see [TrAd]) which are the only superfinitary varieties for which the Kleene theorem holds. (A survey of results connected with varieties is given in [AdTr].)

2. DISTRIBUTIVE LAWS AS NATURAL TRANSFORMATIONS

In this section we present an equivalent characterization of distributive laws using the language of natural transformations between functors.

2.1. THEOREM. *A functor $F: \underline{C} \rightarrow \underline{C}$ satisfies distributive laws over a monad $\underline{T} = (T, \eta, \mu)$ iff there exists a natural transformation $\lambda: FT \rightarrow TF$ such that*

- (i) $\lambda \circ F\eta = \eta F$
- (ii) $\mu F \circ T\lambda \circ \lambda T = \lambda \circ F\mu$

PROOF.

1. Suppose that F satisfies distributive laws over \underline{T} , and let $\{\lambda_A: FTA \rightarrow TFA \mid A \in \text{obj } \underline{C}\}$ be the system of morphisms from 1.4. Let $f: A \rightarrow B$ be a morphism in \underline{C} . According to 1.1 and (1), (2) from 1.4 we have

$$\begin{aligned}
 \lambda_B \circ FTf &= \lambda_B \circ F\mu_B \circ FT\eta_B \circ FTf = \lambda_B \circ F((\eta_B \circ f)^+) = \\
 &= (\lambda_B \circ F(\eta_B \circ f))^+ \circ \lambda_A = \mu_{FB} \circ T\lambda_B \circ TF\eta_B \circ TFf \circ \lambda_A = \\
 &= \mu_{FB} \circ T\eta_{FB} \circ TFf \circ \lambda_A = TFf \circ \lambda_A.
 \end{aligned}$$

Hence, the diagram

$$\begin{array}{ccc}
 FTA & \xrightarrow{\lambda_A} & TFA \\
 FTf \downarrow & & \downarrow TFf \\
 FTB & \xrightarrow{\lambda_B} & TFB
 \end{array}$$

commutes for any morphism $f: A \rightarrow B$ and λ is a natural transformation.

2. Denote by id_{TA} the identity map on TA . By the definition of $^+$ -operation, there is $\mu_A = \text{id}_{TA}^+$ and $\mu_{FA} \circ T\lambda_A = (\lambda_A \circ \text{Fid}_{TA})^+$. By (2), we have

$$\mu_{FA} \circ T\lambda_A \circ \lambda_{TA} = (\lambda_A \circ \text{Fid}_{TA})^+ \circ \lambda_{TA} = \lambda_A \circ \text{Fid}_{TA}^+ = \lambda_A \circ F\mu_A.$$

Hence, (ii) is satisfied.

3. On the other hand, suppose that $\lambda: FT \rightarrow TF$ is a natural transformation satisfying (i) and (ii). Since (i) is the first distributive law, it remains to check only the second distributive law. Let $\alpha: A \rightarrow TB$ be a mapping. By (ii), there is $\lambda_B \circ F\mu_B = \mu_{FB} \circ T\lambda_B \circ \lambda_{TB}$. Since λ is a natural transformation, there is $\lambda_{TB} \circ FT\alpha = TF\alpha \circ \lambda_A$. Therefore,

$$\begin{aligned} (\lambda_B \circ F\alpha)^+ \circ \lambda_A &= \mu_{FB} \circ T\lambda_B \circ TF\alpha \circ \lambda_A = \mu_{FB} \circ T\lambda_B \circ \lambda_{TB} \circ FT\alpha = \\ &= \lambda_B \circ F\mu_B \circ FT\alpha = \lambda_B \circ F\alpha^+. \end{aligned}$$

Hence, (2) is satisfied. \square

2.2. COROLLARY. Let \underline{C} be a category, $F: \underline{C} \rightarrow \underline{C}$, $T: \underline{C} \rightarrow \underline{C}$ be functors, $\underline{T} = (T, \eta, \mu)$ a monad. Then F has an extension to $\underline{C}(\underline{T})$ iff there exists a natural transformation $\lambda: FT \rightarrow TF$ satisfying (i) and (ii) for Theorem 2.1.

3. PROJECTIVE MONADS

In this section we introduce the concept of projective monad and prove that every functor has an extension to the Kleisli category of a projective monad.

3.1. DEFINITION. A monad $\underline{T} = (T, \eta, \mu)$ is called *projective* if there exists a natural transformation $\pi: T \rightarrow \text{Id}$ such that $\pi \circ \eta$ is an identity transformation and $\pi \circ \mu = \pi \circ \pi T$. (We call such a transformation π a projection.)

3.2. REMARKS. 1. For any projective monad (T, η, μ) there is $T\emptyset = \emptyset$.

2. Since π is a natural transformation there is

$$\pi \circ T\pi = \pi \circ \pi T.$$

3.3. EXAMPLE. Sem (see 1.3) is a projective monad.

PROOF. Define a natural transformation $p: T \rightarrow \text{Id}$ by $p_A(a_1 \dots a_n) = a_1$. For any set A and $a, a_{11}, \dots, a_{1k(1)}, \dots, a_{n1}, \dots, a_{nk(n)} \in A$ there is:

$$p_A i_A(a) = p_A(a) = a$$

$$\begin{aligned} p_A s_A((a_{11} \dots a_{1k(1)}) \dots (a_{n1} \dots a_{nk(n)})) &= \\ &= p_A(a_{11} \dots a_{1k(1)} \dots a_{n1} \dots a_{nk(n)}) = a_{11}, \end{aligned}$$

$$p_A p_{SA}((a_{11} \dots a_{1k(1)}) \dots (a_{n1} \dots a_{nk(n)})) = p_A(a_{11} \dots a_{1k(1)}) = a_{11}.$$

Hence, p is a projection. \square

3.4. EXAMPLE. Let n be a positive integer; $(\text{Hom}(n, -), e^{(n)}, m^{(n)})$ is a monad of n -tuples defined as follows:

$$e_A^{(n)}(a) = (\underbrace{a, \dots, a}_n),$$

$$m_A^{(n)}((a_{11}, \dots, a_{1n}), \dots, (a_{n1}, \dots, a_{nn})) = (a_{11}, a_{22}, \dots, a_{nn}).$$

Define

$$p^{(n)}: \text{Hom}(n, -) \rightarrow \text{Id} \quad \text{by} \quad p_A^{(n)}(a_1, \dots, a_n) = a_1.$$

For any set A and $a, a_{11}, \dots, a_{1n}, \dots, a_{n1}, \dots, a_{nn} \in A$ there is

$$p_A^{(n)} e_A^{(n)}(a) = p_A^{(n)}(\underbrace{a, \dots, a}_n) = a,$$

$$\begin{aligned} p_A^{(n)} m_A^{(n)}((a_{11}, \dots, a_{1n}), \dots, (a_{n1}, \dots, a_{nn})) &= \\ &= p_A^{(n)}(a_{11}, a_{22}, \dots, a_{nn}) = a_{11}, \end{aligned}$$

$$\begin{aligned} p_A^{(n)} p_{\text{Hom}(n, A)}^{(n)}((a_{11}, \dots, a_{1n}), \dots, (a_{n1}, \dots, a_{nn})) &= \\ &= p_A^{(n)}(a_{11}, \dots, a_{1n}) = a_{11}. \end{aligned}$$

Hence, $p^{(n)}$ is a projection. \square

3.5. THEOREM. Let \underline{C} be a category, $T: \underline{C} \rightarrow \underline{C}$ a functor, $\underline{T} = (T, \eta, \mu)$ a projective monad (with the projection π). Then an arbitrary functor $F: \underline{C} \rightarrow \underline{C}$ satisfies distributive laws over \underline{T} .

PROOF. Define a natural transformation $\lambda: FT \rightarrow TF$ by $\lambda = \eta F \circ F\pi$. Since \underline{T} is projective there is $\lambda \circ F\eta = \eta F \circ F\pi \circ F\eta = \eta F$ and (i) from 2.1 holds. Since \underline{T} is a monad there is

$$\mu F \circ T\lambda \circ \lambda T = \mu F \circ T\eta F \circ TF\pi \circ \eta FT \circ F\pi T = TF\pi \circ \eta FT \circ F\pi T.$$

Since η is a natural transformation, there is

$$TF\pi \circ \eta FT \circ F\pi T = \eta F \circ F\pi \circ F\pi T = \eta F \circ F\pi \circ F\mu = \lambda \circ F\mu$$

and (ii) from 2.1 holds as well.

According to 2.1, F satisfies distributive laws over \underline{T} . \square

3.6. THEOREM. Let \underline{C} be a category, $T: \underline{C} \rightarrow \underline{C}$ a functor, $\underline{T} = (T, \eta, \mu)$ a projective monad. Then an arbitrary functor $F: \underline{C} \rightarrow \underline{C}$ has an extension to $\underline{C}(\underline{T})$.

PROOF. follows from 1.5 and 3.5. \square

3.7. COROLLARY. An arbitrary set-functor has an extension to the Kleisli category over Sem , i.e. to the category of free semigroups.

PROOF. follows from 3.3 and 3.6. \square

4. CATEGORY OF FREE MONOIDS

While the previous section has solved the question of extensions of set-functors to the category of free semigroups completely and positively, the problem of extensions of set-functors to the category of free monoids seems to be more difficult. In this section, we summarize the known results in this area.

Throughout this section, N denotes the set of all positive integers, \underline{FM} denotes the category of free monoids (i.e. the Kleisli category over Mon), $X \neq \emptyset$ will be a set and $f: X \rightarrow N$ a mapping.

4.1. THEOREM. *If $f(x) = 1$ for any $x \in X$ then $\coprod_{x \in X} \text{Hom}(f(x), -) \cong - \times X$ has an extension to \underline{FM} .*

PROOF. is given in [ArM]. \square

4.2. THEOREM. *If f is bounded, $\max\{f(x) \mid x \in X\} \geq 2$ then $\coprod_{x \in X} \text{Hom}(f(x), -)$ has no extension to \underline{FM} .*

PROOF. is given [V1]. \square

4.3. THEOREM. *If f is a mapping onto $N \setminus \{1\}$ then $F = \coprod_{x \in X} \text{Hom}(f(x), -)$ has an extension to \underline{FM} .*

PROOF. Let A be a set, $x \in X$, $f(x) = n > 1$. Define $\lambda_{Ax}: \text{Hom}(f(x), MA) \rightarrow MFA$ by

$$\lambda_{Ax}(a_{11} \dots a_{1k(1)}, a_{21} \dots a_{2k(2)}, \dots, a_{n1} \dots a_{nk(n)}) = w$$

where

$$w = (a_{11}, a_{21}, a_{22}, \dots, a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)})(a_{12}, a_{21}, a_{22}, \dots, \\ a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)}) \dots (a_{1k(1)}, a_{21}, a_{22}, \dots, \\ a_{n1}, \dots, a_{nk(n)})$$

if $k(1) + (k(2) + \dots + k(n)) > 0$, $w = \Lambda$ otherwise.

Then for any $x \in X$, $v \in \text{Hom}(f(x), MA)$, put $\lambda_A(v) = \lambda_{Ax}(v)$. We have to check conditions (1), (2) from 1.4.

a) There is

$$\begin{aligned} \lambda_A^F(e_A)(\underbrace{a_1, \dots, a_n}_{\in FA}) &= \lambda_A(\underbrace{a_1, \dots, a_n}_{\in FMA}) = (\underbrace{a_1, \dots, a_n}_{\in MFA}) = \\ &= e_{FA}(a_1, \dots, a_n). \end{aligned}$$

b)

$$\begin{array}{ccc}
 \text{FMA} & \xrightarrow{\lambda_A} & \text{MFA} \\
 \text{F}\alpha^+ \downarrow & & \downarrow (\lambda_B \circ \text{F}\alpha)^+ \\
 \text{FMB} & \xrightarrow{\lambda_B} & \text{MFB}
 \end{array}$$

commutes for any $\alpha: A \rightarrow MB$ because

$$\begin{aligned}
 & (\lambda_B \circ \text{F}\alpha)^+ \lambda_A (a_{11} \dots a_{1k(1)}, a_{21} \dots a_{2k(2)}, \dots, a_{n1} \dots a_{nk(n)}) = \\
 & = (\lambda_B \circ \text{F}\alpha)^+ ((a_{11}, a_{21}, \dots, a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)}) (a_{12}, \\
 & \quad a_{21}, \dots, a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)}) \dots (a_{1k(1)}, a_{21}, \\
 & \quad \dots, a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)})) = \mu_{\text{FB}} \circ M(\lambda_B) (\text{F}\alpha(a_{11}, \\
 & \quad a_{21}, \dots, a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)}), \text{F}\alpha(a_{12}, a_{21}, \dots, \\
 & \quad a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)}), \dots, \text{F}\alpha(a_{1k(1)}, a_{21}, \dots, \\
 & \quad a_{2k(2)}, \dots, a_{n1}, \dots, a_{nk(n)})) = \\
 & = \mu_{\text{FB}} \circ M(\lambda_B) ((b_{111} \dots b_{11r(1,1)} \ b_{211} \dots b_{21r(2,1)}, \dots, \\
 & \quad b_{nk(n)1} \dots b_{nk(n)r(n,k(n))}) (b_{121} \dots b_{12r(1,2)}, \\
 & \quad b_{211} \dots b_{21r(2,1)} \dots, b_{nk(n)1} \dots b_{nk(n)r(n,k(n))}) \dots \\
 & \quad \dots (b_{1k(1)1} \dots b_{1k(1)r(1,k(1))}, b_{211} \dots b_{21r(2,1)}, \dots, \\
 & \quad b_{nk(n)1} \dots b_{nk(n)r(n,k(n))})) = \\
 & = (b_{111}, b_{211}, b_{212}, \dots, b_{21r(2,1)}, \dots, b_{nk(n)1}, \dots, \\
 & \quad b_{nk(n)r(n,k(n))}) (b_{112}, b_{211}, b_{212}, \dots, b_{21r(2,1)}, \dots, \\
 & \quad b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))}) \dots (b_{11r(1,1)}, b_{211},
 \end{aligned}$$

$$\begin{aligned}
& b_{212}, \dots, b_{21r(2,1)}, \dots, b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))}) \dots \\
& \dots (b_{1k(1)r(1,k(1))}, b_{211}, b_{212}, \dots, b_{21r(2,1)}, \\
& \dots, b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))})
\end{aligned}$$

where $\alpha(a_{ij}) = b_{ij1} \dots b_{ijr(i,j)}$, and

$$\begin{aligned}
& \lambda_B \text{F}\alpha^+(a_{11} \dots a_{1k(1)}, \dots, a_{n1} \dots a_{nk(n)}) = \\
& = \lambda_B (b_{111} b_{112} \dots b_{1k(1)r(1,k(1))}, \\
& \quad b_{211} b_{212} \dots b_{2k(2)r(2,k(2))}, \dots, \\
& \quad b_{n11} b_{n12} \dots b_{nk(n)r(n,k(n))}) = \\
& = (b_{111}, b_{211}, b_{212}, \dots, b_{21r(2,1)}, \dots, \\
& \quad b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))}) \\
& \quad (b_{112}, b_{211}, b_{212}, \dots, b_{21r(2,1)}, \dots, \\
& \quad b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))}) \dots \\
& \quad \dots (b_{11r(1,1)}, b_{211}, b_{212}, \dots, b_{21r(2,1)}, \dots, \\
& \quad b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))}) \dots \\
& \quad \dots (b_{1k(1)r(1,k(1))}, b_{211}, b_{212}, \dots, \\
& \quad b_{21r(2,1)}, \dots, b_{nk(n)1}, \dots, b_{nk(n)r(n,k(n))}) = \\
& = (\lambda_B \circ \text{F}\alpha)^+ \lambda_A (a_{11} \dots a_{1k(1)}, \dots, a_{n1} \dots a_{nk(n)}).
\end{aligned}$$

Obviously,

$$\begin{aligned}
(\lambda_B \circ F\alpha)^+ \lambda_A(a_{11} \dots a_{1k(1)}, \Lambda, \dots, \Lambda) &= \Lambda = \\
&= \lambda_B^{F\alpha^+}(a_{11} \dots a_{1k(1)}, \Lambda, \dots, \Lambda), \\
(\lambda_B \circ F\alpha)^+ \lambda_A(\Lambda, a_{21} \dots a_{2k(2)}, \dots, a_{n1} \dots a_{nk(n)}) &= \Lambda = \\
&= \lambda_B^{F\alpha^+}(\Lambda, a_{21} \dots a_{2k(2)}, \dots, a_{n1} \dots a_{nk(n)}), \\
(\lambda_B \circ F\alpha)^+ \lambda_A(\Lambda, \dots, \Lambda) &= \Lambda = \lambda_B^{F\alpha^+}(\Lambda, \dots, \Lambda).
\end{aligned}$$

This finishes the proof. \square

4.4. THEOREM. *If f is a mapping onto N then $\bigsqcup_{x \in X} \text{Hom}(f(x), -)$ has an extension to FM.*

PROOF. Denote $Y = \{x \in X \mid f(x) = 1\}$, $Z = \{x \in X \mid f(x) > 1\}$. By Theorem 4.1, $\bigsqcup_{x \in Y} \text{Hom}(f(x), -)$ has an extension to FM. By Theorem 4.3, $\bigsqcup_{x \in Z} \text{Hom}(f(x), -)$ has an extension to FM. Hence, also $\bigsqcup_{x \in X} \text{Hom}(f(x), -)$ has an extension to FM. \square

4.5. THEOREM. $\bigsqcup_{x \in X} \text{Sym}(f(x), -)$ has an extension to FM if either f is a mapping onto N , or $f(x) = 1$ for any $x \in X$.

PROOF. is given in [V2].

5. REFERENCES

- [Ad] J. ADÁMEK, *Automata and categories: finiteness contra minimality*, Math. Found. of Comp. Sci., 4th Symp., Mariánské Lázně^V, Lecture Notes in Comp. Sci. 32 (1975), 160-167.
- [AdTr] J. ADÁMEK and V. TRNKOVÁ, *Varietors and machines in a category*, Algebra Univ. 13 (1981), 89-132.
- [ArM] M.A. ARBIB and E.G. MANES, *Fuzzy machines in a category*, Bull. Austr. Math. Soc. 13 (1975), 169-210.
- [Be] J. BECK, *Distributive laws*, Lecture Notes in Math., Springer 80 (1969), 119-140.
- [BuH] L. BUDACH and H.-J. HOEHNKE, *Automaten und Funktoren*, Akademie-Verlag Berlin 1975.
- [G] J.A. GOGUEN, *Minimal realization of machines in closed categories*, Bull. Amer. Math. Soc. 78 (1972), 777-783.
- [ML] S. MacLANE, *Categories for the working mathematician*, Springer 1972.
- [M] E.G. MANES, *Algebraic theories*, Graduate Texts in Math., Springer 1976.
- [Th] W. THOLEN, *General machines and concrete functors*, Proc. Fund. Comp. Theory 1979, 450-461.
- [TrAd] V. TRNKOVÁ and J. ADÁMEK, *Tree group automata*, Fund. Comp. Theory 1979, 462-467.
- [Tr1] V. TRNKOVÁ, *Automata and categories*, Math. Found. of Comp. Sci., 4th Symp., Mariánské Lázně^V, Lecture Notes in Comp. Sci. 32 (1975), 138-152.
- [Tr2] V. TRNKOVÁ, *L-fuzzy functorial automata*, Math. Found. of Comp. Sci. '79, Lecture Notes in Comp. Sci. 74 (1979), 463-473.
- [V1] J. VINÁREK, *On extensions of functors to the Kleisli category*, Comment. Math. Univ. Carolinae 18 (1977), 319-327.
- [V2] J. VINÁREK, *Extensions of symmetric hom-functors to the Kleisli category*, Fund. Comp. Theory 1981, Lect. Notes in Comp. Sci. 117 (1981), 394-399.

ONTVANGEN 2 4 JUNI 1983