


STICHTING
MATHEMATISCH CENTRUM
2e BOERHAAVESTRAAT 49
AMSTERDAM
AFDELING ZUIVERE WISKUNDE

WN 14

On families of equi-uniformly distributed
sequences in compact spaces

by

P.C. Baayen and G. Helmberg



July 1964

The Mathematical Centre at Amsterdam, founded the 11th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications, and is sponsored by the Netherlands Government through the Netherlands Organization for Pure Research (Z.W.O.) and the Central National Council for Applied Scientific Research in the Netherlands (T.N.O.), by the Municipality of Amsterdam and by several industries.

1. Introduction and preliminaries

Let X be a compact Hausdorff space, let μ be a regular, normed Borel measure on X , and let $C(X)$ be the Banach space (under uniform norm $\|f\| = \sup_{x \in X} |f(x)|$) of all complex-valued continuous functions f on X . Identifying measure and corresponding integral on $C(X)$ we shall use the notation $\mu(f) = \int_X f(x) d\mu(x)$. A family $\mathcal{J} = \{(x_{\sigma,n})_{n=1}^{\infty} : \sigma \in S\}$ of sequences in X is called a family of equi- μ -uniformly distributed sequences (in German: "gleichmässig μ -gleichverteilt") if for every $f \in C(X)$ and for every real number $\varepsilon > 0$ there exists an integer $N(f, \varepsilon)$, independent of σ , such that

$$(1) \quad \left| \frac{1}{N} \sum_{n=1}^N f(x_{\sigma,n}) - \mu(f) \right| \leq \varepsilon \quad \text{for all } N \geq N(f, \varepsilon)$$

and for all $\sigma \in S$ (Hlawka [9]).

The question arises as to the possible size of such a family \mathcal{J} , both from the topological and from the measure-theoretic point of view (section 4). This question may be modified by restricting attention to sequences of a special type only such as the tail sequences of a given μ -uniformly distributed sequence (section 5,6) or the sequences formed by the successive powers of a generator in a compact monothetic group (section 2,3). In all of these modifications, a common feature of the answers to the above questions is the following: such a family \mathcal{J} will in general be in some (topological) sense nowhere dense, it may be enlarged to one which is in some (topological) sense closed and if the set of all sequences in question in some natural sense is given the measure 1, then \mathcal{J} may have measure arbitrary close to 1.

Before taking up the subject in detail we mention two facts which we shall have to use. First, in order to show that the sequences of a certain family \mathcal{J} are equi- μ -

uniformly distributed it obviously suffices to establish (1) for the functions f of a fixed set $F \subset C(X)$ having the property that finite linear combinations of elements of F are uniformly dense in $C(X)$ ([9] § 2, [11] Satz 4). This fact will in the following be referred to as Weyl's criterion.

Secondly we note that, in the definition given above, $C(X)$ may be replaced by the set of all real-valued Borel measurable functions f on X having the following property: for every $\eta > 0$ there exist real-valued continuous functions f_1, f_2 such that $f_1 \leq f \leq f_2$ and $\mu(f_2 - f_1) \leq \eta$. Indeed, we then have

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^N f_1(x_{\sigma,n}) - \mu(f_1) - \eta &\leq \frac{1}{N} \sum_{n=1}^N f_1(x_{\sigma,n}) - \\ &- \mu(f_2) \leq \frac{1}{N} \sum_{n=1}^N f(x_{\sigma,n}) - \mu(f) < \frac{1}{N} \sum_{n=1}^N f_2(x_{\sigma,n}) - \\ &- \mu(f_1) \leq \frac{1}{N} \sum_{n=1}^N f_2(x_{\sigma,n}) - \mu(f_2) + \eta \end{aligned}$$

and therefore

$$\begin{aligned} (2) \quad \left| \frac{1}{N} \sum_{n=1}^N f(x_{\sigma,n}) - \mu(f) \right| &\leq \\ &\leq \max \left\{ \left| \frac{1}{N} \sum_{n=1}^N f_1(x_{\sigma,n}) - \mu(f_1) \right|, \right. \\ &\quad \left. \left| \frac{1}{N} \sum_{n=1}^N f_2(x_{\sigma,n}) - \mu(f_2) \right| \right\} + \eta. \end{aligned}$$

Since η is arbitrary it is indeed possible to choose $N(f, \varepsilon)$ so large that for all $N \geq N(f, \varepsilon)$ the right hand member is smaller than a given $\varepsilon > 0$, uniformly in σ .

As a consequence, if $\mathcal{J} = \{(x_{\sigma,n})_{n=1}^{\infty} : \sigma \in S\}$ is a family of equi- μ -uniformly distributed sequences, (1)

will hold for any complex-valued (Borel measurable) function f whose discontinuities are contained in a closed μ -zero set, in particular for every characteristic function χ_A of a subset A of X whose boundary has μ -measure zero and for the product of χ_A with any function $f \in C(X)$.

A case of particular interest, already mentioned, is the set of sequences $(a^n)_{n=1}^\infty$ if X is a compact monothetic group and if a runs through the generators of X . Here, because of the one-to-one correspondence between generators and sequences, the answers to the questions referred to above may be given by statements about sets of generators of X . Since it is both instructive and convenient for our later considerations, we shall first deal with the special case of uniform distribution mod 1 before passing on to more general situations.

Some of the results of sections 2 and 3 have been announced by the authors in a talk given in October 1963 in the colloquium on uniform distribution at the Mathematical Center in Amsterdam.

2. Uniform distribution mod 1

Let X be the additive group of reals mod 1 and let μ be ordinary Lebesgue measure on $[0, 1[$. It is well known that the sequence $(na)_{n=1}^{\infty}$ is $(\mu -)$ uniformly distributed mod 1 iff a is irrational. Taking in Weyl's criterion for F the set of functions $f_k(x) = e^{2\pi i kx}$ (k an integer) we find that necessary and sufficient for the sequences $(x_{\sigma, n})_{n=1}^{\infty}$ ($\sigma \in S$) to be equi-uniformly distributed mod 1 is the existence, for every integer $k \neq 0$ and for every real number $\epsilon > 0$, of an integer $N(k, \epsilon)$, independent of σ , such that

$$(3) \quad \left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i k x_{\sigma, n}} \right| \leq \epsilon \quad \text{for all } N \geq N(k, \epsilon)$$

and for all $\sigma \in S$. In what follows we shall denote by $[a]$ the greatest integer in a and by $\hat{a} = a - [a]$ the residue of a mod 1. This should not cause confusion with the further use of brackets when replacing parentheses or signalling open or close intervals.

Theorem 1. Let A be a set of irrational numbers such that the sequences $(na)_{n=1}^{\infty}$ ($a \in A$) are equi-uniformly distributed mod 1. Then the set of residues mod 1, $\hat{A} = \{a : a \in A\}$ is nowhere dense in $[0, 1[$.

Proof. Assume the contrary. Then we can find an integer $k \neq 0$ such that the set kA is dense in $[0, 1[\pmod{1}$. Given any integer $N \neq 0$ and any real number δ ($0 < \delta < 1$) we can therefore find a number $a \in A$ such that $ka \in [0, \frac{\delta}{N}] \pmod{1}$ and, a fortiori, $nka \in [0, \delta[$ for $1 \leq n \leq N$. Thus, for these n , if δ has been chosen small enough, $e^{2\pi i nka}$ is close to 1 and so is $\left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i nka} \right|$, in contradiction to (3).

Corollary 1.1. Let a be irrational. Then the sequences $(nma)_{n=1}^{\infty}$ ($m=1, 2, \dots$) are not equi-uniformly distributed mod 1.

Corollary 1.2. The family of all sequences $(na)_{n=1}^{\infty}$

(a irrational) is not the union of countably many families of mod 1 equi-uniformly distributed sequences.

Corollary 1.3. Let A and \hat{A} be as in theorem 1. Then \hat{A} has outer Lebesgue measure smaller than 1.

Theorem 2. Let δ ($0 < \delta < 1$) be given. Then there exists a closed nowhere dense set A of irrational numbers in $[0, 1[$ such that $\mu(A) \geq 1 - \delta$ and such that the sequences $(na)_{n=1}^{\infty}$ ($a \in A$) are equi-uniformly distributed mod 1.

Proof. For every integer $k \geq 1$ we choose a real number $c_k > 0$ in such a way that

$$\mu \{ x \in [0, 1[: |e^{2\pi i k x} - 1| < c_k \} \leq \frac{\delta}{2^k}.$$

Let $A_k = \{ x \in [0, 1[: |e^{2\pi i k x} - 1| \geq c_k \}$ and let

$A = \bigcap_{k=1}^{\infty} A_k$. Then A is closed, consists of irrational numbers only, and we have $\mu(A) \geq 1 - \sum_{k=1}^{\infty} \frac{\delta}{2^k} = 1 - \delta$.

Furthermore, for any $k \geq 1$ and for any $a \in A$ we have

$$\left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i n k a} \right| = \frac{1}{N} \left| \frac{e^{2\pi i N k a} - 1}{e^{2\pi i k a} - 1} \right| \leq \frac{2}{N c_k}.$$

Thus, by Weyl's criterion, the sequences $(na)_{n=1}^{\infty}$ ($a \in A$) are equi-uniformly distributed mod 1 and A is nowhere dense by theorem 1.

3. Uniform distribution in compact monothetic groups

Let X be a compact group which, as a topological space, is Hausdorff, and let μ be normed Haar measure on X . The group X is called monothetic if there is at least one element $a \in X$ (called "generator") such that the sequence $(a^n)_{n=1}^{\infty}$ is everywhere dense in X (van Dantzig [1] cf. [3], [8] II § 9). As a consequence, a compact monothetic group is abelian. If a is a generator (and only in this case), the sequence $(a^n)_{n=1}^{\infty}$ is μ -uniformly distributed [2]. If $(x_n)_{n=1}^{\infty}$ is any μ -uniformly distributed sequence in X , then the sequences $(x_n y) (y \in X)$ are equi- μ -uniformly distributed [9]. We denote by \bar{A} the closure of a subset A of X . In the rest of this section, we shall omit explicit reference to Haar measure μ in statements concerning uniform distribution.

Theorem 3. Let A be a set of generators of X such that the sequences $(a^n)_{n=1}^{\infty} (a \in A)$ are equi-uniformly distributed. Then also the sequences $(a^n)_{n=1}^{\infty} (a \in \bar{A})$ are equi-uniformly distributed.

Proof. Let $f \in C(X)$ and $\epsilon > 0$ be given. Suppose we have

$$(4) \quad \left| \frac{1}{N} \sum_{n=1}^N f(a^n) - \mu(f) \right| \leq \epsilon \quad \text{for all } N \geq N(f, \epsilon)$$

and for all $a \in A$. Then, since $f(x^n)$ is a continuous function of x for every n , (4) also holds for all $a \in \bar{A}$.

We note that theorem 3 yields another proof of theorem 1 which may be regarded as a corollary of theorem 3. Indeed, if A satisfies the hypotheses of theorem 1 and if \bar{A} contained an open interval, then \bar{A} would have to contain rational numbers which certainly are no generators. We may generalize the statement of theorem 1 in the following form:

Theorem 4. Let X be not totally disconnected and let A be a set of generators of X such that the sequences $(a^n)_{n=1}^{\infty} (a \in A)$ are equi-uniformly distributed. Then A is nowhere

dense.

Proof. By Pontrjagin's duality theorem ([16] theorem 46, [8] theorem 24.26) there exists a continuous character d of X such that $d(X) = \{d(x) : x \in X\}$ is the entire circle group T . By theorem 3, without loss of generality we may assume A to be closed. If A contained an open subset of X the $d(A)$ would have to contain an open interval in T since d is an open mapping on X onto T . On the other hand, since d is a continuous homomorphism, $d(A)$ would have to consist of generators of T only, a contradiction.

Corollary 4.1. Suppose that X is connected and a is a generator of X . Then the sequences $((a^m)^n)_{n=1}^\infty$ ($m=1,2,\dots$) are uniformly distributed but not equi-uniformly distributed.

Proof. Since X is connected iff its dual group is torsion-free ([16] theorem 46, [8] theorem 24.25), we have $d(a^m) = d^m(a) \neq 1$ for every non-trivial character d of X and for every integer $m \geq 1$. Thus, a^m is a generator of X ([8] theorem 25.11). The second assertion follows from theorem 4 since the sequence $(a^m)_{m=1}^\infty$ is everywhere dense in X .

We note that, by theorem 4, the second assertion of Corollary 4.1 even holds if X is not totally disconnected, but in this case a^m need not any more be a generator for every $m \geq 2$. For convenience of notation we shall denote the set of all generators of X by G .

Corollary 4.2. Suppose that X is connected and satisfies the second axiom of countability. Then the family of all sequences $(a^n)_{n=1}^\infty$ ($a \in G$) is not the union of countably many families of equi-uniformly distributed sequences.

Proof. Under the stated hypothesis, G is a set of first category in X ([3], cf. [8] 25.27).

Corollary 4.3. Let X and A be as in theorem 4. Then

$$\mu(\bar{A}) < 1.$$

Proof. If $\mu(\bar{A}) = 1$, then \bar{A} would have to be everywhere dense in X since every open set has positive Haar measure.

If X is totally disconnected, the assertion of theorem 4 may fail to hold. In fact, the sequences $(a^n)_{n=1}^{\infty}$ ($a \in G$) will be equi-uniformly distributed (see theorem 5 below) and G may be open (see example following corollary 5.1 below). In order to show this we first establish an auxiliary result. We note that, in the absence of the second axiom of countability, Zorn's lemma is used in the proof.

Lemma 1. Let d be a continuous character of X having the property that $d(X) = T$. Then, for every irrational number α there exists a generator $a \in G$ such that $d(a) = e^{2\pi i \alpha}$.

Proof. Let $D = \{d_\lambda : \lambda \in \Lambda\}$ be a complete system of different continuous characters of X , let d_0 be the trivial character ($d_0 \equiv 1$), and let $d_1 = d$. Since X is monothetic there is at least one generator $b \in G$. Suppose $d_\lambda(b) = e^{2\pi i \beta_\lambda}$ ($\beta_0 = 1$). By assumption ($d_1(X) = T$), β_1 is irrational. Let T_Λ be the topological direct product $T_\Lambda = \prod_{\lambda \in \Lambda} T_\lambda$ ($T_\lambda = T$ for all $\lambda \in \Lambda$). It is well known that the mapping $\varphi: x \rightarrow \hat{x} = (d_\lambda(x))_{\lambda \in \Lambda} \in T_\Lambda$ is a topological isomorphism of X onto a closed subgroup $\hat{X} = \varphi(X)$ of T_Λ . In fact, \hat{X} is the closure (with respect to the product topology in T_Λ) of the sequence $(\hat{b}^n)_{n=1}^{\infty} \subset T_\Lambda$. (The notation is chosen for technical reasons and should not be confused with the notation for residues mod 1 in section 2).

We now choose a maximal system of exponents β_λ , linearly independent over the field of rational numbers and including $\beta_0 = 1$ and β_1 . We denote by $D' = \{d_{\lambda'} : \lambda' \in \Lambda'\}$ the corresponding subsystem of D . We now define an element a of T_Λ in the following way: first we choose arbitrarily a set of real numbers $\alpha_{\lambda'}$ ($\lambda' \in \Lambda'$), linearly independent over the field of rational numbers and including $\alpha_0 = 1$ and $\alpha_1 = \alpha$ (this may even be done by changing at most two of the numbers $\beta_{\lambda'}$). Then we define α_λ for $\lambda \in \Lambda \setminus \Lambda'$ by writing β_λ as a finite linear combination of $\beta_{\lambda'}$'s with rational coefficients and replacing every $\beta_{\lambda'}$

by α_λ . Now we put $\hat{a} = (e^{2\pi i \alpha_\lambda})_{\lambda \in \Lambda} \in T_\Lambda$.

If $U(\hat{a}; \lambda_1, \dots, \lambda_m; \varepsilon) = \{\hat{x} = (e^{2\pi i \xi_\lambda})_{\lambda \in \Lambda} \in T_\Lambda : |e^{2\pi i \xi_\lambda} - e^{2\pi i \alpha_\lambda}| < \varepsilon, k=1, \dots, m\}$ is an arbitrary

neighbourhood of \hat{a} in T_Λ , then we can find an integer n (not necessarily positive) such that $\hat{b}^n \in U(\hat{a}; \lambda_1, \dots, \lambda_m; \varepsilon)$. Indeed, because of our construction of α_λ , for any lattice

point $(h_{\lambda_k})_{k=1}^m$, the congruence $\sum_{k=1}^m h_{\lambda_k} \beta_{\lambda_k} \equiv 0 \pmod{1}$

implies $\sum_{k=1}^m h_{\lambda_k} \alpha_{\lambda_k} \equiv 0 \pmod{1}$.

Thus, by Kronecker's theorem ([12] VII § 2) the simultaneous inequalities

$$|n \beta_{\lambda_k} - \alpha_{\lambda_k}| < \varepsilon \pmod{1} \quad (k=1, \dots, m)$$

have a common integral solution n for any choice of

$\varepsilon = \varepsilon(\varepsilon)$. Thus \hat{a} belongs to the group \hat{X} and therefore there exists an element $a \in X$ such that $\hat{a} = \varphi(a)$.

By our construction α_λ will be an integer if and only if β_λ is an integer. Since b was a generator of X , this is the case if and only if $\lambda=0$. Thus, $d_\lambda(a) \neq 1$ for all $\lambda \neq 0$ and a is a generator of X ([8] theorem 25.11).

Theorem 5. The sequences $(a^n)_{n=1}^\infty$ ($a \in G$) are equi-uniformly distributed if and only if X is totally disconnected.

Proof. We apply Weyl's criterion, taking in place of F a complete system D of different continuous characters of X . If X is totally disconnected, then $d(X)$ is finite for every $d \in D$. For every $a \in G$ and for every non-trivial character $d \in D$, we therefore have $|d(a)-1| \geq c_d > 0$, where c_d only depends on d . Thus, for every non-trivial character d , we obtain

$$\begin{aligned} (5) \quad & \left| \frac{1}{N} \sum_{n=1}^N d(a^n) \right| = \\ & = \frac{1}{N} \left| \frac{d(a^{N+1}) - d(a)}{d(a) - 1} \right| \leq \frac{2}{N c_d} \leq \varepsilon \text{ for all } N \geq \frac{2}{c_d \varepsilon} \end{aligned}$$

and for all $a \in G$.

If X is not totally disconnected, then there is a character $d \in D$ such that $d(X) = T$. If the sequences $(a^n)_{n=1}^\infty$ ($a \in G$) were equi-uniformly distributed, then, by theorem 3, we would have $G = \bar{G}$. In consequence, by lemma 1, we would have $d(G) = T$. This, however, is impossible since $e^{2\pi i \alpha}$ is not a generator of T for rational α .

Corollary 5.1. G is closed if and only if X is totally disconnected.

Proof. If X is totally disconnected, the assertion follows from theorem 5 and theorem 3. If X is not totally disconnected, the same reasoning applies as in the second part of the proof of theorem 5.

Corollary 5.1 may of course also be proved without any reference to equi-uniformly distributed sequences. As a consequence of lemma 1, G can certainly not be open unless X is totally disconnected. If, however, this is the case, then G may well be open as already demonstrated by any finite group with the discrete topology. Below we shall exhibit a compact totally disconnected monothetic group X containing infinitely many elements and having an open set of generators. This, in connection with theorem 5, shows that we cannot omit the hypothesis of X not being totally disconnected in theorem 4.

Let T_∞ be the topological direct product of countably many copies of T . Let $a \in T_\infty$ be defined by $a = (e^{2\pi i \cdot 2^{-k}})_{k=1}^\infty$ and define X to be the closure of the sequence $(a^n)_{n=1}^\infty$ in T_∞ . Then X is a compact monothetic group. Consider an element $b = (e^{2\pi i b_k \cdot 2^{-k}})_{k=1}^\infty \in X$ (b_k integers, $1 \leq b_k \leq 2^k$) with the property that $b_1 = 1$. Since b may be approximated arbitrarily close by elements of the form a^n ($n \geq 1$), for every given $m \geq 1$ the system of simultaneous congruences

and $n \equiv b_k \pmod{2^k} \quad (k=1, \dots, m)$

has a solution $n > 0$ which has to be an odd integer. Therefore, there exists an integer $n' > 0$ such that

$$1 \equiv n' b_k \pmod{2^k} \quad (k=1, \dots, m)$$

and a may be approximated arbitrarily closely by elements of the form $b^{n'}$. Thus, b is a generator and we have

$G = \{ b = (e^{2\pi i b_k \cdot 2^{-k}})_{k=1}^{\infty} : b_1 = 1 \}$. This set G , however, is obviously an open subset of X of Haar measure $\frac{1}{2}$.

The following theorem establishes a generalization of theorem 2 which we, however, shall use in the proof.

We note that, if X satisfies the second axiom of countability, G is a G_δ -set and thus a Borel set ([8] 25.27).

Theorem 6. Suppose that X satisfies the second axiom of countability and let ε ($0 < \varepsilon < 1$) be given. Then there is a closed nowhere dense subset Q of G such that $\mu(Q) > \mu(G) - \varepsilon$ and such that the sequences $(a^n)_{n=1}^{\infty}$ ($a \in Q$) are equi-uniformly distributed.

Proof. Let $D = \{d_k : k=0, 1, 2, \dots\}$ be a complete system of different continuous characters of X and let D' be the subset of D consisting of those characters d for which $d(X) = T$ (D' is empty iff X is totally disconnected). We observe that, λ denoting normed Haar measure on T , for every $d \in D'$ and for every Borel set $B \subset T$, we have $\lambda(B) = \mu(d^{-1}(B))$. (This may be seen e.g. by defining a Borel measure λ on T by this equation and checking that it has all the properties characterizing Haar measure on T). For every character $d_k \in D'$ we may choose by theorem 2 a closed subset A_k of T having the property that the sequences $(d_k(a^n))_{n=1}^{\infty}$ ($d_k(a) \in A_k$) are equi- λ -uniformly distributed in T and such that $\lambda(A_k) > 1 - \frac{\varepsilon}{2^k}$. We put

$Q_1 = \bigcap_{d_k \in D'} d_k^{-1}(A_k)$ if D' is not empty and $Q_1 = X$ if D' is empty. Then $\mu(Q_1) > 1 - \varepsilon$ and, putting $Q = Q_1 \cap G$, we have $\mu(Q) > \mu(G) - \varepsilon$.

Suppose now that $a \in Q$ is chosen arbitrarily and that $d_k \in D$ is any non-trivial character. If $d_k(X) = T$, then $d_k(a) \in A_k$ and

$$(6) \quad \left| \frac{1}{N} \sum_{n=1}^N d_k(a^n) \right| = \left| \frac{1}{N} \sum_{n=1}^N (d_k(a))^n \right| \leq \varepsilon$$

for all $N \geq N(d_k, \varepsilon)$

and for all $a \in Q$. If $d_k(X)$ is finite, then $d_k(a) \neq 1$ since a is a generator and (6) holds by the reasoning used in the proof of (5). Thus, by Weyl's criterion, the sequences $(a^n)_{n=1}^{\infty}$ ($a \in Q$) are equi-uniformly distributed in X . By theorem 3 we may replace Q by its closure which will then have the properties asserted in theorem 6.

We remark that, while no assumption of connectedness is necessary in the hypothesis, theorem 6 is interesting only in the non-totally disconnected case in view of theorem 5.

Corollary 6.1. Suppose that X is connected and satisfies the second axiom of countability. Then, for every given

ε ($0 < \varepsilon < 1$) there is a closed nowhere dense subset A of X such that $\mu(A) > 1 - \varepsilon$ and such that the sequences $(a^n)_{n=1}^{\infty}$ ($a \in A$) are equi-uniformly distributed.

Proof. Under the mentioned hypotheses we have $\mu(G) = 1$ ([3], cf. '8 25.27). An application of theorem 6 completes the proof.

4. Uniform distribution in compact Hausdorff spaces

Let X be any compact Hausdorff space and let μ be any regular normed Borel measure on X . Every sequence $\xi = (x_n)_{n=1}^{\infty} \subset X$ may be regarded as an element of the topological direct product space X_{∞} of countably many copies of X . We denote by $\overline{\mathcal{A}}$ the closure of a subset \mathcal{A} of X_{∞} in the product topology. The theorems stated in the previous sections have rather close analoga also in the present context.

Theorem 7. Suppose that $\mathcal{J} = \{ \xi_{\sigma} \in X_{\infty} : \sigma \in S \}$ is a family of equi-uniformly distributed sequences. Then so is $\overline{\mathcal{J}}$.

Proof: Let $f \in C(X)$ and $\epsilon > 0$ be given. By assumption, we have

$$(7) \quad \left| \frac{1}{N} \sum_{n=1}^N f(x_{\sigma, n}) - \mu(f) \right| \leq \epsilon \quad \text{for all } N \geq N(f, \epsilon)$$

and for all $\sigma \in S$. Let $\xi = (x_n)_{n=1}^{\infty}$ be an arbitrary element of X_{∞} and define $F_N(\xi) = \frac{1}{N} \sum_{n=1}^N f(x_n)$. Then F_N is a continuous function on X_{∞} . Thus, for $\xi \in \mathcal{J}$, we conclude from (7) that also

$$\left| \frac{1}{N} \sum_{n=1}^N f(x_n) - \mu(f) \right| \leq \epsilon \quad \text{for all } N \geq N(f, \epsilon).$$

This proves the assertion.

Theorem 8. Suppose that X contains at least two points and that \mathcal{J} is a family of equi- μ -uniformly distributed sequences. Then $\overline{\mathcal{J}}$ is nowhere dense in X .

Proof. Assume the contrary. Because of theorem 7 we may assume that \mathcal{J} is closed and therefore contains an open subset \mathcal{U} of X . If X satisfies the second axiom of countability, then, via Baire's category theorem, this contradicts the well known fact that the set of all μ -uniformly distributed sequences in X is of first category in X_{∞} .

If X does not satisfy the second axiom of countability we may argue the following way: Without loss of generality we may assume that \mathcal{U} is of the form $\mathcal{U} = \prod_{n=1}^{\infty} U_n$, where U_n is an open subset of X for all $n \geq 1$ and $U_n = X$ for all $n \geq N_1$. Let V and W be open subsets of X with disjoint closures. Let f be a Urysohn function on X (i.e. $f \in C(X)$ and $0 \leq f(x) \leq 1$ for all $x \in X$) such that $f(x) = 1$ for all $x \in V$ and $f(x) = 0$ for all $x \in W$. Let $N \geq 3N_1$ be given. Because of our assumption \mathcal{F} contains sequences

$y = (y_n)_{n=1}^{\infty} \in \mathcal{U}$ and $z = (z_n)_{n=1}^{\infty} \in \mathcal{U}$ with the following properties: $y_n \in V$ for $N_1 < n \leq N$ and $z_n \in W$ for $N_1 < n \leq N$. We conclude $\frac{1}{N} \sum_{n=1}^N f(y_n) \geq \frac{2}{3}$ and $\frac{1}{N} \sum_{n=1}^N f(z_n) \leq \frac{1}{3}$. Since N was arbitrary, this contradicts our assumption that \mathcal{F} is a family of equi- μ -uniformly distributed sequences.

If X satisfies the second axiom of countability, then, as mentioned above, the set of all μ -uniformly distributed sequences in X is of first category in X_{∞} . This could suggest the idea that, in contrast to corollary 1.2 and 4.2, the set of all μ -uniformly distributed sequences might well be the countable union of families of equi- μ -uniformly distributed sequences. A closer inspection, however, shows that this still will not be the case in general. Indeed, if X is a compact connected abelian group satisfying the second axiom of countability (and a fortiori monothetic, [8] 25.14) and if μ is normed Haar measure on X , then the set of all μ -uniformly distributed sequences in particular contains all sequences $(a^n)_{n=1}^{\infty}$ ($a \in G$). This set, however, cannot be split into the union of countably many families of equi- μ -uniformly distributed sequences by corollary 4.2.

In order to state an analogon of corollary 4.1 (cf. [7] theorem 2) we recall that the shift transformation P in X_{∞} is defined by $P(x_n)_{n=1}^{\infty} = (y_n)_{n=1}^{\infty}$, where $y_n = x_{n+1}$ ($1 \leq n < \infty$).

Corollary 8.1. Suppose that X contains at least two points. Suppose furthermore that $\xi \in X_\infty$ is μ -uniformly distributed in X and that, for some subsequence $(m_k)_{k=1}^\infty$ of the positive integers, the sequence $(p^{m_k} \xi)_{k=1}^\infty$ is dense in some open subset of X_∞ . Then the sequences $P^{m_k} \xi$ ($k=1,2,\dots$) are μ -uniformly distributed but not equi- μ -uniformly distributed.

Corollary 8.2. Let X and \mathcal{T} be as in theorem 8 and suppose that X is the support of μ . Then $\mu_\infty(\bar{\mathcal{T}}) < 1$.

Proof. Under the mentioned hypothesis, the support of μ_∞ is X_∞ . If $\bar{\mathcal{T}}$ had μ_∞ -measure 1, then $\bar{\mathcal{T}}$ would coincide with X_∞ , in contradiction to theorem 8.

The following theorem and its proof are essentially due to E. Hlawka ([9] § 6) who has stated them for the case of a compact group.

Theorem 9. Suppose that X satisfies the second axiom of countability and let δ ($0 < \delta < 1$) be given. Then there exists a closed family \mathcal{T} of equi- μ -uniformly distributed sequences such that $\mu_\infty(\mathcal{T}) > 1 - \delta$.

Proof. In order to be able to apply Weyl's criterion we first note that because of the second axiom of countability there exists a countable set $F = \{f_k: k=1,2,\dots\} \subset C(X)$ as described in section 1. Given any $k \geq 1$ we again define $F_{k,N} \in C(X_\infty)$ for $\xi = (x_n)_{n=1}^\infty$ by

$$F_{k,N}(\xi) = \frac{1}{N} \sum_{n=1}^N f_k(x_n).$$

It is a well known consequence of the individual ergodic theorem ([5], [9] § 6, [11]) that

$$\lim_{N \rightarrow \infty} F_{k,N}(\xi) = \mu(f_k)$$

for μ_∞ -almost all $\xi \in X_\infty$. Therefore, by Egoroff's theorem ([4] § 21 A), there exists a measurable subset \mathcal{T}_k of X_∞ of measure $\mu_\infty(\mathcal{T}_k) \geq 1 - \frac{\delta}{k}$ such that the functions $F_{k,N}$ converge uniformly on \mathcal{T}_k for $N \rightarrow \infty$.

Taking $\mathcal{J}_0 = \bigcap_{k=1}^{\infty} \mathcal{J}_k$ we obtain $\mu_{\infty}(\mathcal{J}_0) \geq 1 - \delta$. Furthermore, for every $k \geq 1$ and for every given $\varepsilon > 0$, we have

$$\left| \frac{1}{N} \sum_{n=1}^N f_k(x_n) - \mu(f_k) \right| \leq \varepsilon \quad \text{for all } N \geq N(f_k, \varepsilon)$$

and for all $\xi = (x_n)_{n=1}^{\infty} \in \mathcal{J} = \bar{\mathcal{J}}_0$. An application of Weyl's criterion leads to the assertion.

5. Almost well distributed sequences

Let X be a compact Hausdorff space and let μ be a regular normed Borel measure on X . As in the preceeding section we denote by P the shift transformation in the infinite product space X_∞ . A sequence $\xi \in X_\infty$ is called μ -well distributed if the sequences $P^m \xi$ ($m=1,2,\dots$) are equi- μ -uniformly distributed (Hlawka [9], Petersen [15]). While under the assumption of the second axiom of countability μ_∞ -almost every sequence is μ -uniformly distributed ([11]), μ_∞ -almost no sequence is μ -well distributed if μ is not a point-measure ([7] corollary 2.1). We shall now apply the results of the preceeding section to some questions connected with the concept of good distribution.

Theorem 10. Suppose that the sequence $\xi \in X_\infty$ is μ -well distributed. Then so is every sequence in the closure of the set $\mathcal{F} = \{P^m \xi : m=1,2,\dots\}$.

Proof. By theorem 7, \mathcal{F} is a family of equi- μ -uniformly distributed sequences. If η belongs to \mathcal{F} , then so does $P^m \eta$ for every $m \geq 1$.

As an illustration in the case of uniform distribution mod 1 we note that, if a is irrational and if ξ is the sequence $(na)_{n=1}^\infty \pmod{1}$, then \mathcal{F} is the family of all sequences $(na+b)_{n=1}^\infty \pmod{1}$ ($b \in [0,1[$).

We shall call a sequence $\xi \in X_\infty$ almost μ -well distributed if there exists an infinite subsequence $(m_k)_{k=1}^\infty$ of the non-negative integers such that the sequences $P^{m_k} \xi$ ($k=1,2,\dots$) are equi- μ -uniformly distributed. If we want to refer to this particular subsequence we shall call ξ almost μ -well distributed $(m_k)_{k=1}^\infty$. Note that the very definition implies that an almost μ -well distributed sequence is μ -uniformly distributed.

Theorem 11. Suppose that $\xi \in X_\infty$ is almost μ -well distributed $(m_k)_{k=1}^\infty$ and let h be any positive integer. Then ξ is also almost μ -well distributed $(m_k+h)_{k=1}^\infty$.

Proof. For any given $f \in C(X)$ and for any $\varepsilon > 0$ we have

$$\left| \frac{1}{N} \sum_{n=m_k+1}^{m_k+N} f(x_n) - \mu(f) \right| \leq \varepsilon \quad \text{for all } N \geq N(f, \varepsilon)$$

and for all $k \geq 1$. We conclude

$$\begin{aligned} & \left| \frac{1}{N} \sum_{n=m_k+h+1}^{m_k+h+N} f(x_n) - \mu(f) \right| \leq \\ & \leq \frac{1}{N} \left| \sum_{n=m_k+N+1}^{m_k+N+h} f(x_n) - \sum_{n=m_k+1}^{m_k+h} f(x_n) \right| + \\ & + \left| \frac{1}{N} \sum_{n=m_k+1}^{m_k+N} f(x_n) - \mu(f) \right| \leq \\ & \leq \frac{2h \|f\|}{N} + \varepsilon \leq 2\varepsilon \quad \text{for all } N \geq \max(N(f, \varepsilon), \frac{2h \|f\|}{\varepsilon}) \end{aligned}$$

and for all $k \geq 1$.

Corollary 11.1. Let ξ and h be as in theorem 11. Then the sequences $P_{m_k+m}^{\xi}$ ($|m| \leq h, k=1,2,\dots$) are equi- μ -uniformly distributed.

Proof. By theorem 11, for every fixed m , the sequences $P_{m_k+m}^{\xi}$ ($k=1,2,\dots$) are equi- μ -uniformly distributed.

Taking the union of these finitely many families of sequences, we again obtain a family of equi- μ -uniformly distributed sequences.

Corollary 11.2. Suppose that $\xi \in X_\infty$ is almost μ -well distributed $(m_k)_{k=1}^\infty$ and that the differences $(m_{k+1}-m_k)$ are bounded. Then ξ is μ -well distributed.

Proof. Put $m_0=0$, let $h = \max_{k=0} (m_{k+1}-m_k)$ and apply corollary 11.1.

If we compare these statements with the results of the previous sections, we may regard corollary 11.1 as some sort of analogon to theorem 3 and theorem 7. In fact, it asserts that, restricting attention to the subsequence $P^m \xi$ of a given μ -uniformly distributed sequence ξ , every $m_k \xi$ family of equi- μ -uniformly distributed subsequences $P^{m_k} \xi$ ($k=1,2,\dots$) may be enlarged by an operation similar to taking the closure, referring, however, not to a topology but to the concept of relative density of a set of integers. In this sense, corollary 11.2 appears to be an analogon to theorem 4 and theorem 8 if we state it in the following form: Suppose that $\xi \in X_\infty$ is not μ -well distributed and suppose that the sequences $P^{m_k} \xi$ ($k=1,2,\dots$) are equi- μ -uniformly distributed. Then the sequence of integers $(m_k)_{k=1}^\infty$ is not relatively dense (i.e. it is not possible to find a positive integer h such that every interval of length h contains at least one integer m_k).

The following theorem again stems from a remark of E. Hlawka ([9] last paragraph; the statements "jede Folge" and "offene Menge" in the passage referred to are erroneous). In the context just mentioned, it establishes an analogon to theorem 6 and theorem 9.

Theorem 12. Suppose that X satisfies the second axiom of countability. Then μ_∞ -almost every sequence $\xi \in X_\infty$ is almost μ -well distributed. In particular, μ_∞ -almost every sequence $\xi \in X_\infty$ has the following property: given any δ ($0 < \delta < 1$) there exists a subsequence $(m_k)_{k=1}^\infty$ of the non-negative integers of asymptotic density greater than $1 - \delta$ such that ξ is almost μ -well distributed $(m_k)_{k=1}^\infty$.

Proof. By theorem 9 we can find a sequence of families \mathcal{X}_j of equi- μ -uniformly distributed sequences such that $\mu_\infty(\mathcal{X}_j) \geq 1 - \frac{1}{j}$ ($j=1,2,\dots$). We note that P is a μ_∞ -measure preserving ergodic transformation in X_∞ [5].

By the individual ergodic theorem, there is a subset Z_j

of X such that $\mu_\infty(\gamma_j) = 1$ and such that for every $\xi \in \gamma_j$ the sequence of exponents m_k for which $P_{\infty, \xi}^{m_k} \in \gamma_j$ has asymptotic density $\mu_\infty(\gamma_j)$. Taking $\gamma = \bigcap_{j=1}^{\infty} \gamma_j$ we obtain $\mu_\infty(\gamma) = 1$ and every $\xi \in \gamma$ has the property stated in the theorem.

6. Almost well, weakly well, and completely uniformly distributed sequences

Let, as in section 5, X be a compact Hausdorff space and let μ be a regular normed Borel measure on X . In view of theorem 12 we are led to consider the following classes of sequences in X and to ask for the set-theoretic and measure-theoretic relationships between them:

$\mathcal{U} = \{\xi \in X_\infty : \xi \text{ is } \mu\text{-uniformly distributed}\}.$

$\mathcal{AW} = \{\xi \in X_\infty : \xi \text{ is almost } \mu\text{-well distributed}\}.$

$\mathcal{AW}_1 = \{\xi \in X_\infty : \text{for every } \delta (0 < \delta < 1) \text{ there exists a subsequence } (m_k)_{k=1}^\infty \text{ of the non-negative integers of asymptotic density greater than } 1-\delta \text{ such that } \xi \text{ is almost } \mu\text{-well distributed } (m_k)_{k=1}^\infty\}.$

$\mathcal{AW}_2 = \{\xi \in X_\infty : \text{there exists a subsequence } (m_k)_{k=1}^\infty \text{ of the non-negative integers of asymptotic density } 1 \text{ such that } \xi \text{ is almost } \mu\text{-well distributed } (m_k)_{k=1}^\infty\}.$

$\mathcal{W} = \{\xi \in X : \xi \text{ is } \mu\text{-well distributed}\}.$

Obviously we have $\mathcal{U} \supset \mathcal{AW} \supset \mathcal{AW}_1 \supset \mathcal{AW}_2 \supset \mathcal{W}$. Furthermore, if the second axiom of countability is satisfied, we have $\mu_\infty(\mathcal{AW}_1) = 1$ by theorem 12 and $\mu_\infty(\mathcal{W}) = 0$ if μ is not a point measure by [7] corollary 2.1. In what follows we shall show that, under these hypotheses and in the presence of μ -well distributed sequences, each of the above inclusions is strict and $\mu_\infty(\mathcal{AW}_2) = 0$.

In this context it seems appropriate also to clarify the relations between the properties of a sequence $\xi \in X_\infty$ to be almost μ -well distributed, weakly μ -well distributed in the sense of Hlawka [10] and completely μ -uniformly distributed in the sense of Korobov [13]. We shall therefore also consider the following classes of sequences in X :

$\mathcal{WAW} = \{\xi \in X_\infty : \xi \text{ is weakly } \mu\text{-well distributed, i.e. for every } f \in C(X) \text{ we have}$

$$\lim_{N \rightarrow \infty} \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| = 0 \}.$$

$\mathcal{CU} = \{ \xi \in X_\infty : \xi \text{ is completely } \mu\text{-uniformly distributed, i.e. the sequence } (P^m \xi)_{m=1}^\infty \text{ is } \mu_\infty\text{-uniformly distributed in } X_\infty \}.$

Again, it is well known and easy to see that we have $\mathcal{U} \supset \mathcal{CU}$ and $\mathcal{U} \supset \mathcal{WU} \supset \mathcal{WU}_1$. Furthermore, if the second axiom of countability is satisfied, we have $\mu_\infty(\mathcal{CU}) = 1$ (replacing in the corresponding statement about uniformly distributed sequences X by X_∞ and μ by μ_∞), $\mu_\infty(\mathcal{WU}) = 1$ by [10] Satz 2 (cf. theorem 16), and $\mathcal{CU} \cap \mathcal{WU} = \emptyset$ by [7] theorem 2 if μ is not a point measure. Thus, μ_∞ -almost every sequence $\xi \in X_\infty$ is at the same time almost μ -well distributed (even in the sense of \mathcal{WU}_1), weakly μ -well distributed, and completely μ -uniformly distributed. We shall show that, in fact, $\mathcal{CU} \cap \mathcal{WU}_2 = \emptyset$ if μ is not a point measure, thus establishing $\mu_\infty(\mathcal{WU}_2) = 0$, and that, in the presence of μ -well distributed sequences, all inclusions in the chain $\mathcal{U} \supset \mathcal{WU} \supset \mathcal{WU}_1$ are strict. In particular, we have $\mathcal{WU} \supset \mathcal{WU}_2$ but $\mathcal{WU} \not\supset \mathcal{WU}_1$. Part of the proofs will be carried out by explicitly constructing sequences that are of one type but not of the other.

Let us note again that, in the definition of weakly μ -well distributed sequences, we can replace $\mathcal{C}(X)$ by the larger class of functions described at the end of section 1. In fact, if $\xi \in X_\infty$ is weakly μ -well distributed and if f, f_1, f_2 are real-valued Borel measurable functions ($f_j \in \mathcal{C}(X)$, $j=1,2$) satisfying $f_1 \leq f \leq f_2$ and $\mu(f_2 - f_1) \leq \frac{\varepsilon}{2}$, then we can choose $N(f, \varepsilon)$ so large that for all $N \geq N(f, \varepsilon)$ we have

$$\limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f_j(x_n) - \mu(f_j) \right| \leq \frac{\varepsilon}{4} \quad (j=1,2).$$

By (2) we conclude

$$\begin{aligned}
& \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| \leq \\
& \leq \sum_{j=1}^2 \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f_j(x_n) - \mu(f_j) \right| + \frac{\varepsilon}{2} \leq \varepsilon.
\end{aligned}$$

This proves our assertion.

We now proceed with the detailed exposition of the statements outlined above. The proof of the following theorem is, with some slight modifications, the same as the proof of theorem 2 in [7]. We shall, however, reproduce it here for the convenience of the reader.

Theorem 13. Suppose that μ is not a point measure and suppose that $\xi = (x_n)_{n=1}^{\infty} \in X_{\infty}$ is completely μ -uniformly distributed. Then ξ is not almost μ -well distributed with respect to any subsequence $(m_k)_{k=1}^{\infty}$ of the non-negative integers of asymptotic density 1.

Proof. Assume the contrary and let $M = (m_k)_{k=1}^{\infty}$ be a subsequence of the non-negative integers of asymptotic density 1 such that ξ is almost μ -well distributed $(m_k)_{k=1}^{\infty}$. Since μ is not concentrated in one point we can find an open set $A \subset X$ such that $0 < \mu(A) < 1$. Without loss of generality we may assume that the boundary of A has μ -measure zero. Let $N \geq 1$ be given and let $A_{\infty} = \bigcap_{n=1}^{\infty} A_n$ where $A_n = A$ for $1 \leq n \leq N$ and $A_n = X$ for $n > N$. Then A_{∞} is open in X_{∞} and its boundary has μ_{∞} -measure zero. Furthermore, we have $0 < \mu_{\infty}(A) < 1$. Since the sequence $(P^m \xi)_{m=1}^{\infty}$ is by assumption μ_{∞} -uniformly distributed in X_{∞} and, since M has asymptotic density 1, so is the sequence $(P^{m_k} \xi)_{k=1}^{\infty}$. Therefore, there exists a positive integer $k(N)$ such that $P^{m_{k(N)}} \xi \in A_{\infty}$. Hence, for every choice of $N \geq 1$, we have

$$\frac{1}{N} \sum_{n=m_{k(N)}+1}^{m_{k(N)}+N} \chi_A(x_n) - \mu(A) = 1 - \mu(A) > 0.$$

This, however, contradicts our assumption that ξ is almost μ -well distributed $(m_k)_{k=1}^\infty$.

Corollary 13.1. Suppose that X satisfies the second axiom of countability and that μ is not a point measure. Then $\mu_\infty(\mathcal{AN}_2) = 0$.

Proof. Under the mentioned hypothesis, we have $\mu_\infty(\mathcal{CN}) = 1$. The statement then follows from theorem 13.

Corollary 13.2. Suppose that X satisfies the second axiom of countability and that μ is not a point measure. Then $\mathcal{AN}_1 \not\subset \mathcal{AN}_2$.

Proof. The statement follows from theorem 12 and corollary 13.1.

Theorem 14. Suppose that X contains at least two points and that there exists a μ -well distributed sequence. Then there also exists a sequence $\xi \in X_\infty$, constructed explicitly below, which is not μ -well distributed but almost μ -well distributed $(m_k)_{k=1}^\infty$ where $(m_k)_{k=1}^\infty$ has asymptotic density 1. Consequently we have $\mathcal{AN}_2 \not\subset \mathcal{AN}$.

Proof. Suppose that the sequence $\eta = (y_k)_{k=1}^\infty \in X_\infty$ is μ -well distributed and let $a \in X$ belong to the support of μ . We then choose a point $b \neq a$ in X and construct a sequence $\xi = (x_n)_{n=1}^\infty \in X_\infty$ by induction as follows: Let $x_1 = b$. If x_1, \dots, x_{h^4} ($h \geq 1$ an integer) have already been defined such as to comprise all elements y_k with $k \leq h^4 - h(2h-1)$, then let x_n ($h^4 < n \leq h^4 + 4h^3 + 6h^2$) be the next $4h^3 + 6h^2$ consecutive y_k 's and let $x_n = b$ for $h^4 + 4h^3 + 6h^2 < n \leq (h+1)^4$.

Obviously, due to the increasing stretches of b 's which certainly do not cover the support of μ , the sequence ξ is not μ -well distributed. If we allow m to run through all integers satisfying $h^4 \leq m < h^4 + 4h^3$ ($h=1, 2, \dots$), then it is easy to see that this subsequence \mathcal{M} of the positive integers has asymptotic density 1. We shall now show that the corresponding sequences $P^m \xi$ ($m \in \mathcal{M}$) are equi- μ -uniformly distributed.

To this purpose, let $f \in C(X)$ ($f \not\equiv 0$) and $\varepsilon > 0$ be given

and suppose that

$$\left| \frac{1}{N} \sum_{n=m+1}^{m+N} f(y_n) - \mu(f) \right| \leq \frac{\varepsilon}{2} \quad \text{for all } N \geq N(f, \varepsilon)$$

and for all $m=1, 2, \dots$. It will suffice to show that

$$(8) \quad \left| \frac{1}{N} \sum_{n=m+1}^{m+N} f(x_n) - \mu(f) \right| \leq \varepsilon \quad \text{for all } N \geq N(f, \varepsilon)$$

and for all $m \in \mathbb{N}$ which are greater than the fourth power of some fixed integer $H=H(f, \varepsilon)$. In fact, we choose this integer H in such a way that $\frac{4H+1}{6H^2+4H+1} < \frac{\varepsilon}{4\|f\|}$ and $6H^2 \geq N(f, \varepsilon)$.

For a fixed $m \in \mathbb{N}$, $m \geq H^4$, let us denote by N' the number of elements x_{m+n} ($1 \leq n \leq N$) chosen from the sequence η and by $\sum' f(x_n)$ the sum over these elements. For all $N \geq N(f, \varepsilon)$ we have $N' \geq N(f, \varepsilon)$ because of $H^4 \leq m < H^4 + 4H^3$ and $6H^2 \geq N(f, \varepsilon)$. Thus we obtain

$$\begin{aligned} \frac{1}{N} \sum_{n=m+1}^{m+N} f(x_n) &= \frac{N'}{N} \cdot \frac{1}{N'} \sum' f(x_n) + \frac{N-N'}{N} f(b) = \\ &= \frac{1}{N} \sum' f(x_n) - \frac{N-N'}{N} \cdot \frac{1}{N'} \sum' f(x_n) + \frac{N-N'}{N} f(b) \end{aligned}$$

and

$$\begin{aligned} (9) \quad \left| \frac{1}{N} \sum_{n=m+1}^{m+N} f(x_n) - \mu(f) \right| &\leq \left| \frac{1}{N} \sum' f(x_n) - \mu(f) \right| + 2 \frac{N-N'}{N} \|f\| \leq \\ &\leq \frac{\varepsilon}{2} + 2 \frac{N-N'}{N} \|f\|. \end{aligned}$$

Because of our construction of ξ and of the restriction imposed on m we have

$$\frac{N-N'}{N} \leq \frac{4H+1}{6H^2+4H+1} \leq \frac{4H+1}{6H^2+4H+1} \leq \frac{\varepsilon}{4\|f\|}.$$

This, in connection with (9), proves (8).

In order to establish the inclusion $\mathcal{M} \supset \mathcal{M}_2$ we again need a lemma.

Lemma 2. Let \mathcal{M} be a subsequence of the non-negative integers of asymptotic density 1 and let N be given. Then the subsequence $\mathcal{R} = \{r: rN \in \mathcal{M}\}$ of the non-negative integers also has asymptotic density 1.

Proof. For any integer $M \geq 1$ we define

$$v(\mathcal{M}, M) = \sum_{\substack{m \in \mathcal{M} \\ m < M}} 1.$$

(for other sequences, the counting function v is defined analogously). Suppose $\liminf_{R \rightarrow \infty} \frac{v(\mathcal{R}, R)}{R} = 1 - 2\delta$ ($\delta > 0$), contrary to the assertion. We then have

$$v(\mathcal{M}, RN) \leq v(\mathcal{R}, R) + (N-1)R \quad \text{for every } R \geq 1$$

and therefore

$$\frac{v(\mathcal{M}, RN)}{RN} = \frac{1}{N} \cdot \frac{v(\mathcal{R}, R)}{R} + \frac{N-1}{N} < \frac{1}{N}(1-\delta) + \frac{N-1}{N} = 1 - \frac{\delta}{N}$$

for infinitely many R , in contradiction to our assumption.

Theorem 15. Suppose that the sequence $\xi = (x_n)_{n=1}^{\infty} \in X_{\infty}$ is almost μ -well distributed $(m_k)_{k=1}^{\infty}$ and $\mathcal{M} = (m_k)_{k=1}^{\infty}$ has asymptotic density 1. Then ξ is weakly μ -well distributed.

Consequently, we have $\mathcal{M} \supset \mathcal{M}_2$.

Proof. Let $f \in C(X)$ and $\varepsilon > 0$ be given and let

$$\left| \frac{1}{N} \sum_{n=m+1}^{m+N} f(x_n) - \mu(f) \right| \leq \varepsilon \quad \text{for all } N \geq N(f, \varepsilon)$$

and for all $m \in \mathcal{M}$. Fix any $N \geq N(f, \varepsilon)$ and let $\mathcal{R} = \{r \geq 0: rN \in \mathcal{M}\}$. Furthermore, let \mathcal{R}' be the subsequence of the non-negative integers, complementary to \mathcal{R} . Then we have

$$a_{N,r} = \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| \leq \varepsilon \quad \text{for all } r \in \mathcal{R}$$

and $a_{N,r} \leq 2\|f\|$ for all $r \geq 0$. Since \mathcal{R} has asymptotic density 1 by lemma 2, \mathcal{R}' has asymptotic density 0. We conclude

$$\begin{aligned} \frac{1}{R} \sum_{r=0}^{R-1} a_{N,r} &= \frac{1}{R} \left(\sum_{\substack{r \in \mathcal{R} \\ r < R}} a_{N,r} + \sum_{\substack{r \in \mathcal{R}' \\ r < R}} a_{N,r} \right) \leq \\ &< \frac{R\varepsilon}{R} + \frac{v(\mathcal{R}', R)}{R} \cdot 2\|f\| \leq 2\varepsilon \quad \text{for all } R \geq R_0. \end{aligned}$$

As a consequence, we have

$$\limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} a_{N,r} \leq 2, \quad \text{for all } N \geq N(f, \varepsilon)$$

and, since ε was arbitrary,

$$\lim_{N \rightarrow \infty} \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| = 0.$$

For the case of a compact group X with normed Haar measure μ , the following theorem has been proved by Hlawka ([10] Satz 2). In order to obtain an even sharper statement, the proof given there actually makes use of group theoretic concepts. We shall therefore give another proof here which makes use of ergodic theory and is valid also in the more general situation considered in this section (the possibility of this approach has also already been mentioned by Hlawka).

Theorem 16. Suppose that X satisfies the second axiom of countability. Then μ_∞ -almost all sequences $\xi \in X_\infty$ are weakly μ -well distributed.

Proof. We note that Weyl's criterion holds for weakly μ -well distributed sequences just as well. We therefore choose a countable subset $F = \{f_k : k=1, 2, \dots\} \subset C(X)$ as described in section 1 and fix a function $f = f_k \in F$ ($f \neq 0$). We then define the function $F_N \in C(X_\infty)$ for $\xi = (x_n)_{n=1}^\infty$ by

$$F_N(\xi) = \left| \frac{1}{N} \sum_{n=1}^N f(x_n) - \mu(f) \right|.$$

Since μ_∞ -almost all $\xi \in X_\infty$ are μ -uniformly distributed, we have $\lim_{N \rightarrow \infty} F_N(\xi) = 0$ for μ_∞ -almost all $\xi \in X_\infty$.

Let the integer $j \geq 1$ be given. By Egoroff's theorem ([4] §21 A) there exists a subset \mathcal{J}_j of X_∞ such that $\mu_\infty(\mathcal{J}_j) > 1 - \frac{1}{4j\|f\|}$ and such that the functions F_N converge to zero uniformly on \mathcal{J}_j . Let $N(j)$ be chosen in such a way that $F_N(\xi) \leq \frac{1}{2j}$ for all $N \geq N(j)$ and for all $\xi \in \mathcal{J}_j$. We fix an integer $N \geq N(j)$. By the individual ergodic theorem, applied to the measure preserving transformation P^N in X_∞ , for all ξ in some subset $(f, j, N) \in X_\infty$ of μ_∞ -measure 1, we have $p^{rN}\xi \in \mathcal{J}_j$ for all $r \geq 0$ except for a subsequence $(r_h)_{h=1}^\infty$ of the non-negative integers of asymptotic density smaller than $\frac{1}{4j\|f\|}$. For every such $\xi = (x_n)_{n=1}^\infty \in \mathcal{J}(f, j, N)$ we obtain

$$\begin{aligned} \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| &= \\ &= \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} F_N(p^{rN}\xi) \leq \\ &\leq \limsup_{R \rightarrow \infty} \left(\frac{1}{R} \sum_{\substack{r=0 \\ p^{rN}\xi \in \mathcal{J}_j}}^{R-1} F_N(p^{rN}\xi) + \frac{1}{R} \sum_{\substack{r=0 \\ p^{rN}\xi \notin \mathcal{J}_j}}^{R-1} 2\|f\| \right) \leq \\ &\leq \frac{1}{2j} + \frac{2\|f\|}{4j\|f\|} = \frac{1}{j}. \end{aligned}$$

For every $\xi \in \bigcap_{N=N(j)} \mathcal{J}(f, j, N)$ we therefore have

$$\limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| \leq \frac{1}{j} \quad \text{for all } N \geq N(j).$$

and for every $\xi \in \mathcal{Z}(f) = \bigcap_{j=1}^{\infty} \bigcap_{N=N(j)}^{\infty} \mathcal{Z}(f, j, N)$ we have

$$(10) \quad \lim_{N \rightarrow \infty} \limsup_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} f(x_n) - \mu(f) \right| = 0.$$

Taking $\mathcal{Z} = \bigcap_{k=1}^{\infty} \mathcal{Z}(f_k)$ we finally have $\mu_{\infty}(\mathcal{Z}) = 1$ and, by Weyl's criterion, (10) holds for all $f \in C(X)$ and for all $\xi = (x_n)_{n=1}^{\infty} \in \mathcal{Z}$.

Corollary 16.1. Suppose that X satisfies the second axiom of countability and that μ is not a point measure. Then $\mathcal{W} \supsetneq \mathcal{W}_2$.

Proof. The statement follows from theorem 16 and corollary 13.1.

In view of the inclusions $\mathcal{W} \supsetneq \mathcal{W}_2$, theorem 14 furnishes an example of a sequence which is weakly μ -well distributed but not μ -well distributed. Our last two theorems will be concerned with constructions of sequences belonging to the (differences of) classes $\mathcal{W} \setminus (\mathcal{W}_1 \cup \mathcal{W}_2)$ and $\mathcal{W}_1 \setminus (\mathcal{W}_1 \cup \mathcal{W}_2)$ respectively.

Lemma 3. Let A be closed subset of X with μ -zero boundary and such that $\mu(A) = \alpha > 0$. If the sequence $\xi = (x_n)_{n=1}^{\infty}$ is μ -uniformly (resp. μ -well) distributed in X , then the subsequence $\eta = \xi \cap A = (y_k)_{k=1}^{\infty}$, consisting of all elements of ξ lying in A , is $\frac{1}{\alpha}\mu$ -uniformly (resp. $\frac{1}{\alpha}\mu$ -well) distributed in A , considered as a compact Hausdorff space in the relative topology.

Proof. Since every continuous complex-valued function on A is of the form $f|_A$ ($f \in C(X)$) by Tietze's extension theorem, we have to show that

$$(11) \quad \lim_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{k=1}^N f|_A(y_k) - \frac{1}{\alpha} \mu(f|_A) \right| = 0 \quad \text{for all } f \in C(X),$$

resp. that

$$\lim_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{k=m+1}^{m+N} f\chi_A(y_k) - \frac{1}{\alpha} \mu(f\chi_A) \right| = 0 \quad \text{for all } f \in C(X)$$

uniformly in $m=0,1,2,\dots$

Let $f \in C(X)$ be given and suppose first that ξ is μ -uniformly distributed in X . For every given integer $N \geq 1$, we define N' to be the smallest integer having the property

$$\text{that } \sum_{n=1}^{N'} \chi_A(x_n) = N.$$

Thus, we have $N' \geq N$ and

$$\lim_{N \rightarrow \infty} \frac{N}{\alpha N'} = \frac{1}{\alpha} \cdot \lim_{N' \rightarrow \infty} \frac{1}{N'} \sum_{n=1}^{N'} \chi_A(x_n) = 1.$$

From the identity

$$\left| \frac{1}{N} \sum_{k=1}^N f\chi_A(y_k) - \frac{1}{\alpha} \mu(f\chi_A) \right| = \frac{1}{\alpha} \left| \frac{\alpha N'}{N} \cdot \frac{1}{N'} \sum_{n=1}^{N'} f\chi_A(x_n) - \mu(f\chi_A) \right|$$

we then deduce (11).

Suppose next that ξ is μ -well distributed. For every integer $m \geq 0$ we now define m' as the smallest integer having the property that $\sum_{n=1}^{m'+1} \chi_A(x_n) = m+1$, and $N(m)$ as the smallest integer having the property that $\sum_{n=m'+1}^{m'+N(m)} \chi_A(x_n) = N$. Thus, we have $N(m) \geq N$. Furthermore, by the same reasoning as above, if N is greater than a fixed integer N_0 , independent of m , then $\frac{N}{\alpha N(m)}$ is close to 1 and the difference

$$\left| \frac{1}{N} \sum_{k=m+1}^{m+N} f\chi_A(y_k) - \frac{1}{\alpha} \mu(f\chi_A) \right| = \frac{1}{\alpha} \left| \frac{\alpha N(m)}{N} \cdot \frac{1}{N(m)} \sum_{n=m'+1}^{m'+N(m)} f\chi_A(x_n) - \mu(f\chi_A) \right|$$

is close to 0, uniformly in $m=0,1,2,\dots$. This proves the assertion.

Theorem 17. Suppose that μ is not a point measure and that there exists a μ -uniformly distributed sequence. Then

there also exists a sequence $\xi \in X_\infty$, constructed explicitly below, which is μ -uniformly distributed but not almost μ -well distributed and not weakly μ -well distributed. Consequently we have $\mathcal{U} \neq \mathcal{U} \cap \mathcal{W}$ and $\mathcal{U} \neq \mathcal{W} \cap \mathcal{W}$.

Proof. Let A be a closed subset of X with μ -zero boundary and such that $0 < \mu(A) = \alpha < 1$ (our hypothesis on μ implies the existence of such a set A). Furthermore, let B be the closure of the complement of A . Then also B has μ -zero boundary and we have $\mu(f) = \mu(f\chi_A) + \mu(f\chi_B)$ for every $f \in C(X)$. By lemma 3 there exists a sequence $\eta = (y_k)_{k=1}^\infty$ which is $\frac{1}{\alpha}$ - μ -uniformly distributed in A and a sequence $\zeta = (z_l)_{l=1}^\infty$ which is $\frac{1}{1-\alpha}$ - μ -uniformly distributed in B . We construct the sequence $\xi = (x_n)_{n=1}^\infty$ by induction as follows: Let $x_1 = z_1$, $x_2 = z_2$. If $x_1, \dots, x_{2^{(h)}}^{(h)}$ ($h \geq 1$) have already been chosen such as to comprise certain initial segments of the sequences η and ζ , then let $x_{2^{(h)}+1}, \dots, x_{2^{(h)}+2[h\alpha]}$ be the next $2[h\alpha]$ consecutive y_k 's and let $x_{2^{(h)}+2[h\alpha]+1}, \dots, x_{2^{(h+1)}}^{(h+1)}$ be the next $2h-2[h\alpha]$ consecutive z_l 's not yet incorporated into the sequence ξ to be constructed.

We first show that the sequence ξ constructed in this way is μ -uniformly distributed in X . Let $f \in C(X)$ and $\epsilon > 0$ be given. For a given integer $N \geq 2$ we determine the integer h in such a way that $2^{(h)} \leq N < 2^{(h+1)}$. Furthermore, we define N_y (resp. N_z) to be the number of y_k 's (resp. z_l 's) occurring among the first $2^{(h)}$ x_n 's. Thus, we have $N_{yz} = N_y + N_z = 2^{(h)}$ and

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^N f(x_n) - \mu(f) &= \\ &= \frac{N_{yz}}{N} \left\{ \frac{N_y}{N_{yz}} \left[\frac{1}{N_y} \sum_y f(y_k) - \frac{1}{\alpha} \mu(f\chi_A) \right] + \left[\frac{N_y}{N_{yz}} \cdot \frac{1}{\alpha} - 1 \right] \mu(f\chi_A) \right\} + \\ &+ \frac{N_{yz}}{N} \left\{ \frac{N_z}{N_{yz}} \left[\frac{1}{N_z} \sum_z f(z_l) - \frac{1}{1-\alpha} \mu(f\chi_B) \right] + \left[\frac{N_z}{N_{yz}} \cdot \frac{1}{1-\alpha} - 1 \right] \mu(f\chi_B) \right\} + \end{aligned}$$

$$+ \frac{1}{N} \left[\sum_{n=N_{yz}+1}^N f(x_n) - (N-N_{yz})\mu(f) \right];$$

here \sum_y and \sum_z denote the sums over the y_k 's and z_1 's, respectively, occurring among the first N_{yz} x_n 's, and the third term disappears for $N=N_{yz}$. In view of $\lim_{N \rightarrow \infty} \frac{N_y}{N} = \lim_{N \rightarrow \infty} \frac{N_z}{N} = \alpha$ we obtain, for all sufficiently large N ,

$$\begin{aligned} \left| \frac{1}{N} \sum_{n=1}^N f(x_n) - \mu(f) \right| &\leq \frac{N_y}{N} \varepsilon + \left| \frac{1}{\alpha} \cdot \frac{N_y}{N} - 1 \right| \|f\| + \\ &+ \frac{N_z}{N} \varepsilon + \left| \frac{1}{1-\alpha} \cdot \frac{N_z}{N} - 1 \right| \|f\| + \frac{4h}{N} \|f\|. \end{aligned}$$

Evidently, for sufficiently large N , the last term becomes smaller than ε . Furthermore, we have

$$\frac{N_y}{N_{yz}} = \frac{2[\alpha] + 2 \cdot [2\alpha] + 2 \cdot [3\alpha] + \dots + 2[(h-1)\alpha]}{2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot (h-1)}.$$

Therefore, we conclude $\lim_{N \rightarrow \infty} \frac{N_y}{N} = \alpha$ and, by a similar reasoning, $\lim_{N \rightarrow \infty} \frac{N_z}{N} = 1-\alpha$. Thus, we obtain

$$\left| \frac{1}{N} \sum_{n=1}^N f(x_n) - \mu(f) \right| < 4\varepsilon$$

for all sufficiently large N .

We next show that ξ is not almost μ -well distributed. In order to do this, for every given integer $m \geq 2$ we estimate

the difference $\left| \frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) - \alpha \right|$ from below for suitably chosen integers N . We distinguish four cases, depending on the choice of m .

1) $2\binom{h}{2} \leq m < 2\binom{h}{2} + [h\alpha]$. We choose $N = 2\binom{h}{2} + 2[h\alpha] - m$. Then we have $N > [h\alpha]$ and

$$\left| \frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) - \alpha \right| = 1 - \alpha > 0.$$

2) $2\binom{h}{2} + [h\alpha] \leq m < 2\binom{h}{2} + 2[h\alpha]$. We choose $N = 2\binom{h+1}{2} - m$. Then we have $N > 2h - 2[h\alpha]$ and

$$\frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) \leq \max_{1 \leq j \leq h} \frac{j}{2h - 2[h\alpha] + j} = \frac{[h\alpha]}{2h - [h\alpha]} \leq \frac{\alpha}{2 - \alpha} < \alpha.$$

Therefore, we obtain

$$\left| \frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) - \alpha \right| \geq \alpha - \frac{\alpha}{2 - \alpha} = \alpha \cdot \frac{1 - \alpha}{2 - \alpha} > 0.$$

3) $2\binom{h}{2} - 2[h\alpha] \leq m < 2\binom{h}{2} + h + [h\alpha]$. We choose $N = 2\binom{h+1}{2} - m$. Then we have $N > h - [h\alpha]$ and

$$\left| \frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) - \alpha \right| = \alpha > 0.$$

4) $2\binom{h}{2} + h + [h\alpha] \leq m < 2\binom{h+1}{2}$. We choose $N = 2\binom{h+1}{2} + 2[(h+1)\alpha] - m$. Then we have $N > 2[(h+1)\alpha]$ and

$$\frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) \geq \frac{2[(h+1)\alpha]}{h - [h\alpha] + 2[(h+1)\alpha]}.$$

For large h the right member is close to $\frac{2\alpha}{1+\alpha} > \alpha$. As a consequence, for larger h we obtain

$$\left| \frac{1}{N} \sum_{n=m+1}^{m+N} x_A(x_n) - \alpha \right| \geq \frac{1}{2} \left(\frac{2\alpha}{1+\alpha} - \alpha \right) = \frac{\alpha}{2} \cdot \frac{1-\alpha}{1+\alpha} > 0.$$

Choosing now $\varepsilon = \min(\alpha \cdot \frac{1-\alpha}{2-\alpha}, \frac{\alpha}{2} \cdot \frac{1-\alpha}{1+\alpha})$, we see that, given any positive integer N_0 , for every integer m which is larger than a certain constant M depending on N_0 we can find an integer $N_m \geq N_0$ such that

$$\left| \frac{1}{N_m} \sum_{n=m+1}^{m+N_m} x_A(x_n) - \alpha \right| \geq \varepsilon.$$

It is therefore not possible to select an infinite increasing sequences of integers m_k in such a way that the sequence $P^{m_k} \xi$ ($k=1,2,\dots$) are equi- μ -uniformly distributed. Thus, ξ is not almost μ -well distributed.

Finally we show that the sequence ξ is not weakly μ -well distributed. In fact, we shall prove that, for every fixed integer $N \geq 1$,

$$\lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} x_A(x_n) - \alpha \right| = 2\alpha(1-\alpha).$$

To this purpose it will suffice to show the following: Given any integer $h \geq 2$, let $(x_{rN+1}, \dots, x_{(r+1)N})$ ($r_{h-1} < r \leq r_h$) be those blocks of N consecutive elements of ξ that have some overlap with the block of $2[h\alpha]$ consecutive elements y_k and $2h-2[h\alpha]$ consecutive elements z_1 , but no overlap with the following block of $2[(h+1)\alpha]$ consecutive elements y_k and $2(h+1)-2[(h+1)\alpha]$ consecutive elements z_1 in the same sequence ξ . Then

$$\lim_{h \rightarrow \infty} \frac{1}{r_h - r_{h-1}} \sum_{r=r_{h-1}+1}^{r_h} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} x_A(x_n) - \alpha \right| = 2\alpha(1-\alpha).$$

(We note that $\frac{2h}{N} - 1 \leq r_h - r_{h-1} \leq \frac{2h}{N} + 1$. From this follows $\lim_{h \rightarrow \infty} (r_h - r_{h-1}) = \infty$ and

$$\begin{aligned} \lim_{h \rightarrow \infty} \frac{1}{r_h - r_{h-1}} \sum_{r=r_{h-1}+1}^{r_h} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} x_A(x_n) - \alpha \right| &= \\ &= \lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} x_A(x_n) - \alpha \right|. \end{aligned}$$

Now, if h is sufficiently large, for approximately $\frac{2[h\alpha]}{N}$ (at least $\frac{2[h\alpha]}{N} - 2$) values of r ($r_{h-1} < r \leq r_h$) the difference $d_r =$

$\left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} x_A(x_n) - \alpha \right|$ will have the value $1-\alpha$, while for

approximately $\frac{2h-2[h\alpha]}{N}$ (at least $\frac{2h-2[h\alpha]}{N}-2$) values of r ($r_{h-1} < r \leq r_h$) this value will be α and for at most two values of r , d_r will be different from α and $1-\alpha$ but bounded above by 2. Thus we obtain

$$\lim_{h \rightarrow \infty} \frac{1}{r_h - r_{h-1}} \left| \frac{1}{N} \sum_{n=r_{h-1}N+1}^{r_h N} x_A(x_n) - \alpha \right| =$$

$$\lim_{h \rightarrow \infty} \frac{N}{2h} \cdot \frac{2[h\alpha](1-\alpha) + (2h-2[h\alpha])\alpha}{N} = 2\alpha(1-\alpha).$$

This completes the proof.

In the mod 1 case, a less abstract example of a sequence enjoying the properties mentioned in theorem 17 (having also served as a model for the construction carried out in the proof) is furnished by the sequence $(0, 0, \frac{1}{2}, 0, \frac{1}{3}, \frac{2}{3}, 0, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, 0, \frac{1}{5}, \dots)$. Sequences of this type have been considered in [14] in connection with the rearrangement of arbitrary sequences, everywhere dense mod 1, to mod 1 uniformly distributed sequences.

Theorem 18. Suppose that μ is not a point measure and that there exists a μ -well distributed sequence. Then there also exists an almost μ -well but not weakly μ -well distributed sequence $\omega \in X_\infty$, constructed explicitly below, with the following property: if $M = (m_k)_{k=1}^\infty$ is any infinite subsequence of the non-negative integers such that ω is almost μ -well distributed $(m_k)_{k=1}^\infty$, then M has asymptotic density 0. Consequently we have $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \omega_{m_k} = 0$ and $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \omega_{m_k} \neq 0$.

Proof. As in the proof of theorem 17, let A be a closed subset of X with μ -zero boundary and such that $0 < \mu(A) = \alpha < 1$, and let B be the closure of the complement of A . Let $\xi = (x_k)_{k=1}^\infty$ be a sequence which is μ -well distributed in X , let $\eta = (y_l)_{l=1}^\infty$ be $\frac{1}{\alpha}$ - μ -well distributed in A and let $\zeta = (z_m)_{m=1}^\infty$ be $\frac{1}{1-\alpha}$ - μ -well distributed in B (cf. lemma 3). We construct the sequence $\omega = (u_n)_{n=1}^\infty$ by induction as follows: Let $u_1 = x_1$. If u_1, \dots, u_h ($h \geq 1$) have already been chosen such as to

comprise certain initial segments of the sequences ξ , η , and ζ , let $u_{h^4+1}, \dots, u_{h^4+6h^2+1}$ be the next $6h^2+1$ consecutive x_k 's, and let $u_{h^4+6h^2+2}, \dots, u_{(h+1)^4}$ consist of the $2h^2+2$ consecutive blocks of $2h$ elements each (referred to in the following as 'yz-blocks'), which in turn consist of the next $2[h\alpha]$ consecutive y_1 's and, immediately following, the next $2h-2[h\alpha]$ consecutive z_m 's not yet incorporated into the sequence ω to be constructed.

We shall first show that the sequence ω is almost μ -well distributed $(h^4)_{h=1}^\infty$. To this purpose, given any $f \in C(X)$ and any $\varepsilon > 0$, we choose N_1 such that, uniformly in $h=1, 2, \dots$,

$$\left| \frac{1}{N} \sum_{k=h+1}^{h+N} f(x_k) - \mu(f) \right| \leq \varepsilon$$

$$\left| \frac{1}{N} \sum_{l=h+1}^{h+N} f(y_l) - \mu(f\chi_A) \right| \leq \varepsilon \quad \text{for all } N \geq N_1.$$

$$\left| \frac{1}{N} \sum_{m=h+1}^{h+N} f(z_m) - \mu(f\chi_B) \right| \leq \varepsilon$$

Furthermore, we choose the integer H such that $\min(2[H\alpha], 2H-2[H\alpha]) \geq N_1$ and such that $\max\left(\frac{4h}{6h^2+1}, \left|\frac{[h\alpha]}{h\alpha} - 1\right|, \left|\frac{h-[h\alpha]}{h(1-\alpha)} - 1\right|\right) \leq \varepsilon$ for all $h \geq H$. Let now $h \geq H$ and $N \geq N_1$ be given. We define N_x to be the number of x_k 's among the elements $u_{h^4+1}, \dots, u_{h^4+N}$. Furthermore, consider the yz-blocks completely covered by these elements and let N_y and N_z be the number of the y_1 's and z_m 's contained in all of these yz-blocks, respectively. We set $N_{yz} = N_y + N_z$ and $N_u = N - N_x - N_y - N_z$. Then we have

$$\frac{1}{N} \sum_{n=h^4+1}^{h^4+N} f(u_n) - \mu(f) = \frac{N_x}{N} \left[\frac{1}{N_x} \sum_x f(x_k) - \mu(f) \right] +$$

$$\begin{aligned}
& + \frac{N_{yz}}{N} \left\{ \frac{N_y}{N_{yz}} \left[\frac{1}{N_y} \sum_y f(y_1) - \frac{1}{\alpha} \mu(f \chi_A) \right] + \left[\frac{N_y}{\alpha N_{yz}} - 1 \right] \mu(f \chi_A) \right\} + \\
& + \frac{N_{yz}}{N} \left\{ \frac{N_z}{N_{yz}} \left[\frac{1}{N_z} \sum_z f(z_m) - \frac{1}{1-\alpha} \mu(f \chi_B) \right] + \left[\frac{N_z}{(1-\alpha)N_{yz}} - 1 \right] \mu(f \chi_B) \right\} + \\
& + \frac{1}{N} \left[\sum_u f(u_n) - N_u \mu(f) \right].
\end{aligned}$$

Here, the sums \sum_x , \sum_y , \sum_z are extended over the corresponding N_x, N_y, N_z elements x_k, y_1, z_m respectively. The second and third member disappear if $N < 6h^2 + 1 + 2h$; the fourth member also may disappear and contains a sum \sum_u over N_u tail elements where N_u (if positive) is less than the number of elements in the unfinished yz -block. Taking absolute values, because of $N \geq N_1$ and $6h^2 + 1 \geq \min(2[h\alpha], 2h - 2[h\alpha]) \geq N_1$ we obtain

$$\begin{aligned}
\left| \frac{1}{N} \sum_{n=h+1}^{h^4+N} f(u_n) - \mu(f) \right| & \leq \frac{N_x}{N} \varepsilon + \\
& + \frac{N_{yz}}{N} \left\{ \frac{N_y}{N_{yz}} \varepsilon + \left| \frac{N_y}{\alpha N_{yz}} - 1 \right| \|f\| \right\} + \\
& + \frac{N_{yz}}{N} \left\{ \frac{N_z}{N_{yz}} \varepsilon + \left| \frac{N_z}{(1-\alpha)N_{yz}} - 1 \right| \|f\| \right\} + \\
& + \frac{2h}{6h^2+1} \cdot 2\|f\| \leq \\
& \leq \varepsilon + \left| \frac{N_y}{\alpha N_{yz}} - 1 \right| \|f\| + \left| \frac{N_z}{(1-\alpha)N_{yz}} - 1 \right| \|f\| + \varepsilon \|f\|.
\end{aligned}$$

Let the integer $i \geq 1$ be chosen such that $(h+i-1)^4 < h^4 + N \leq (h+i)^4$. We then have

$$\begin{aligned}
N_{yz} &= 2h(2h^2+2) + 2(h+1)(2(h+1)^2+2) + \dots + 2(h+i-1) \cdot j \\
N_y &= 2[h\alpha](2h^2+2) + 2[(h+1)\alpha](2(h+1)^2+2) + \dots + 2[(h+i-1)\alpha] \cdot j \\
N_z &= 2(h-[h\alpha])(2h^2+2) + 2(h+1-[(h+1)\alpha])(2(h+1)^2+2) + \dots + \\
&\quad + 2(h+i-1-[(h+i-1)\alpha]) \cdot j
\end{aligned}$$

and therefore

$$\min\left(\frac{[h\alpha]}{h}, \dots, \frac{[(h+i-1)\alpha]}{h+i-1}\right) \leq \frac{N_{yz}}{N_y} \leq \max\left(\frac{[h\alpha]}{h}, \dots, \frac{[(h+i-1)\alpha]}{h+i-1}\right).$$

By our choice of $h \geq H$ we conclude $\left|\frac{N_y}{\alpha N_{yz}} - 1\right| \leq \varepsilon$ and, analogously, $\left|\frac{N_z}{(1-\alpha)N_{yz}} - 1\right| \leq \varepsilon$. Thus we obtain

$$\left| \frac{1}{N} \sum_{n=h^4+1}^{h^4+N} f(u_n) - \mu(f) \right| \leq \varepsilon + 3\varepsilon \|f\| \quad \text{for all } N \geq N_1$$

and for all $h \geq H$. This proves that ω is almost μ -well distributed $(h^4)_{h=1}^\infty$.

Suppose next that the subsequence $\mathcal{M} = (m_k)_{k=1}^\infty$ of the non-negative integers has positive upper asymptotic density. We shall show that the sequences $P_{m_k} \omega$ ($k=1,2,\dots$) cannot be equi- μ -uniformly distributed. Consider the set of positive integers $\mathcal{N} = \{n: h^4 - 6(h-1)^2 \leq n \leq h^4 + 6h^2 \text{ for some } h=1,2,\dots\}$. Since \mathcal{N} has asymptotic density zero, infinitely many $m_k \in \mathcal{M}$ must lie outside of \mathcal{N} and therefore satisfy the inequality

$$h^4 + 6h^2 + 1 \leq m_k \leq (h+1)^4 - 6h^2 - 1 \quad (h \geq 1 \text{ an integer})$$

It follows that the element u_{m_k+1} belongs to some yz -block of length $2h$ and, since $m_k + 4h < (h+1)^4$, the elements u_{m_k+n} ($1 \leq n \leq 4h$) cover at least one complete yz -block of length $2h$. Since this happens for infinitely many k , by the same reasoning as in the proof of theorem 17, the

sequences $P^m_k \omega$ ($k=1,2,\dots$) cannot be equi- μ -uniformly distributed.

Finally, since the elements x_k form a subsequence of asymptotic density 0 in ω , by the same reasoning as in the proof of theorem 17 we find that

$$\lim_{R \rightarrow \infty} \frac{1}{R} \sum_{r=0}^{R-1} \left| \frac{1}{N} \sum_{n=rN+1}^{(r+1)N} x_A(u_n)^{-\alpha} \right| = 2\alpha(1-\alpha).$$

for all $N \geq 1$. Thus, the sequence ω cannot be weakly μ -well distributed. This completes the proof.

The preceding discussion still leaves some open questions concerning the remaining possible inclusions between the classes $\mathcal{O} \cap \mathcal{W}$ and $\mathcal{O} \cap \mathcal{W}_1$ on the one hand and $\mathcal{W} \cap \mathcal{W}_1$ on the other hand. Apart from these we mention one particular question to which we do not know the answer: Suppose that X satisfies the second axiom of countability. Does there always exist a μ -well distributed sequence?

References

- 1 van Dantzig, D.: Zur topologischen Algebra I. Komplet-
tierungstheorie. Math. Ann. 107, 587-626 (1932).
- 2 Eckmann, B.: Über monothetische Gruppen. Commentarii
math. Helvet. 16, 249-263 (1943/44).
- 3 Halmos, P.R. and Samelson, H.: On monothetic groups.
Proc. Nat. Acad. Sci. USA 28, 254-258 (1942).
- 4 Halmos, P.R.: Measure Theory. Van Nostrand, New York
1956.
- 5 : Lectures on Ergodic Theory. Publications
of the Math. Society of Japan 3, Tokyo 1956.
- 6 Hartman, S. und Ryll-Nardzewski, C.: Zur Theorie der
lokalkompakten abelschen Gruppen. Colloquium
mathematicum 4, 157-188 (1956).
- 7 Helmsberg, G. and Paalman-de Miranda, A.: Almost no
sequence is well distributed. To appear in Proc.
Kon. Ned. Ak. Wetenschappen = Indag. Math.
- 8 Hewitt, E. and Ross, K.A.: Abstract Harmonic Analysis
I. Springer Verlag, Berlin 1963.
- 9 Hlawka, E.: Zur formalen Theorie der Gleichverteilung
in kompakten Gruppen. Rend. Circ. mat. Palermo 4,
33-47 (1955).
- 10 : Über einen Satz von van der Corput. Archiv
der Math. 6, 115-120 (1955).
- 11 : Folgen auf kompakten Räumen. Abh. math.
Sem. Univ. Hamburg 20, 223-241 (1956).
- 12 Koksma, J.F.: Diophantische Approximationen. Ergebn.
Math. u. Grenzgeb. Bd. 4, Heft 4, Springer Verlag,
Berlin 1936.
- 13 Korobov, N.M.: On completely uniform distribution and
jointly normal numbers (Russian). Izvestija Akad.
Nauk SSSR, Ser. Math. 20, 649-660 (1956).

- 14 von Neumann, J.: Gleichmässig dichte Zahlenfolgen
(Hungarian, German summary). Math. fiz. Lapok 32,
32-40 (1925).
- 15 Petersen, G.M.: Almost convergence and uniformly distributed sequences. Quart. J. Math., Oxford, 2. ser.
7, 188-191 (1956).
- 16 Pontrjagin, L.S.: Topologische Gruppen I,II. Teubner
Verlag, Leipzig 1957.

1. The first part of the report is a general introduction to the subject of the study. It discusses the importance of the study and the objectives of the research.

2. The second part of the report is a detailed description of the methodology used in the study. It includes information about the sample, the data collection methods, and the statistical analysis.

3. The third part of the report is a discussion of the results of the study. It compares the findings with the previous research and discusses the implications of the study.