

NAG C Library Chapter Introduction

e01 – Interpolation

Contents

1	Scope of the Chapter	2
2	Background to the Problems	2
3	Recommendations on Choice and Use of Available Functions	3
3.1	General	3
3.2	One Independent Variable	3
3.2.1	Interpolated values: data without derivatives	3
3.2.2	Interpolating function: data without derivatives	3
3.2.3	Data containing derivatives	4
3.3	Two Independent Variables	4
3.3.1	Data on a rectangular mesh	4
3.3.2	Arbitrary data	4
3.4	Three Independent Variables	5
3.4.1	Arbitrary data	5
4	Decision Tree	5
5	Index	5
6	Functions Withdrawn or Scheduled for Withdrawal	6
7	References	6

1 Scope of the Chapter

This chapter is concerned with the interpolation of a function of one, two or three variables. When provided with the value of the function (and possibly one or more of its lowest-order derivatives) at each of a number of values of the variable(s), the functions provide either an interpolating function or an interpolated value. For some of the interpolating functions, there are supporting functions to evaluate, differentiate or integrate them.

2 Background to the Problems

In motivation and in some of its numerical processes, this chapter has much in common with Chapter e02 (Curve and Surface Fitting). For this reason, we shall adopt the same terminology and refer to dependent variable and independent variable(s) instead of function and variable(s). Where there is only one independent variable, we shall denote it by x and the dependent variable by y . Thus, in the basic problem considered in this chapter, we are given a set of distinct values x_1, x_2, \dots, x_m of x and a corresponding set of values y_1, y_2, \dots, y_m of y , and we shall describe the problem as being one of interpolating the data points (x_r, y_r) , rather than interpolating a function. In modern usage, however, **interpolation** can have either of two rather different meanings, both relevant to functions in this chapter. They are

- (a) the determination of a function of x which takes the value y_r at $x = x_r$, for $r = 1, 2, \dots, m$ (an **interpolating function** or **interpolant**),
- (b) the determination of the value (**interpolated value** or **interpolate**) of an interpolating function at any given value, say \hat{x} , of x within the range of the x_r (so as to estimate the value at \hat{x} of the function underlying the data).

The latter is the older meaning, associated particularly with the use of mathematical tables. The term ‘function underlying the data’, like the other terminology described above, is used so as to cover situations additional to those in which the data points have been computed from a known function, as with a mathematical table. In some contexts, the function may be unknown, perhaps representing the dependency of one physical variable on another, say temperature upon time.

Whether the underlying function is known or unknown, the object of interpolation will usually be to approximate it to acceptable accuracy by a function which is easy to evaluate anywhere in some range of interest. Polynomials, rational functions (ratios of two polynomials) and piecewise polynomials, such as cubic splines (see Section 2.2 in the e02 Chapter Introduction for definitions of terms in the latter case), being easy to evaluate and also capable of approximating a wide variety of functions, are the types of function mostly used in this chapter as interpolating functions. An interpolating polynomial is taken to have degree $m - 1$ when there are m data points, and so it is unique. It is called the **Lagrange interpolating polynomial**. The rational function, in the special form used, is also unique. An interpolating spline, on the other hand, depends on the choice made for the knots.

One way of achieving the objective in (b) above is, of course, through (a), but there are also methods which do not involve the explicit computation of the interpolating function. Everett’s formula and Aitken’s successive linear interpolation (see Dahlquist and Björck (1974)) provide two such methods. Both are used in this chapter and determine a value of the Lagrange interpolating polynomial.

It is important to appreciate, however, that the Lagrange interpolating polynomial often exhibits unwanted fluctuations between the data points. These tend to occur particularly towards the ends of the data range, and to get larger with increasing number of data points. In severe cases, such as with 30 or 40 equally spaced values of x , the polynomial can take on values several orders of magnitude larger than the data values. (Closer spacing near the ends of the range tends to improve the situation, and wider spacing tends to make it worse.) Clearly, therefore, the Lagrange polynomial often gives a very poor approximation to the function underlying the data. On the other hand, it can be perfectly satisfactory when its use is restricted to providing interpolated values away from the ends of the data range from a reasonably small number of data values.

In contrast, a cubic spline which interpolates a large number of data points can often be used satisfactorily over the whole of the data range. Unwanted fluctuations can still arise but much less frequently and much less severely than with polynomials. Rational functions, when appropriate, would also be used over the whole data range. The main danger with these functions is that their polynomial denominators may take zero values within that range. Unwanted fluctuations are avoided altogether by a function using piecewise

cubic polynomials having only first derivative continuity. It is designed especially for monotonic data, but for other data still provides an interpolant which increases, or decreases, over the same intervals as the data.

The concept of interpolation can be generalized in a number of ways. Firstly, at each x , the interpolating function may be required to take on not only a given value but also given values for all its derivatives up to some specified order (which can vary with r). This is the Hermite–Birkoff interpolation problem. Secondly, we may be required to estimate the value of the underlying function at a value \hat{x} outside the range of the data. This is the process of **extrapolation**. In general, it is a good deal less accurate than interpolation and is to be avoided whenever possible.

Interpolation can also be extended to the case of two or more independent variables. If the data values are given at the intersections of a regular two-dimensional mesh bicubic splines (see Section 2.3.2 in the e02 Chapter Introduction) are very suitable and usually very effective for the problem. For other cases, perhaps where the data values are quite arbitrarily scattered, polynomials and splines are not at all appropriate and special forms of interpolating function have to be employed. Many such forms have been devised and two of the most successful are in functions in this chapter. They both have continuity in first, but not higher, derivatives.

3 Recommendations on Choice and Use of Available Functions

3.1 General

Before undertaking interpolation, in other than the simplest cases, you should seriously consider the alternative of using a function from Chapter e02 to approximate the data by a polynomial or spline containing significantly fewer coefficients than the corresponding interpolating function. This approach is much less liable to produce unwanted fluctuations and so can often provide a better approximation to the function underlying the data.

When interpolation is employed to approximate either an underlying function or its values, you will need to be satisfied that the accuracy of approximation achieved is adequate. There may be a means for doing this which is particular to the application, or the function used may itself provide a means. In other cases, one possibility is to repeat the interpolation using one or more extra data points, if they are available, or otherwise one or more fewer, and to compare the results. Other possibilities, if it is an interpolating function which is determined, are to examine the function graphically, if that gives sufficient accuracy, or to observe the behaviour of the differences in a finite-difference table, formed from evaluations of the interpolating function at equally-spaced values of x over the range of interest. The spacing should be small enough to cause the typical size of the differences to decrease as the order of difference increases.

3.2 One Independent Variable

3.2.1 Interpolated values: data without derivatives

When the underlying function is well represented by data points on both sides of the value, x , at which an interpolated value is required, `nag_1d_cheb_interp` (e01aec) can be used to construct the interpolating polynomial. In other cases the cubic spline, `nag_1d_spline_interpolant` (e01bac), is more generally applicable and is particularly appropriate when interpolated values towards the end of the range are required. It is also likely to be preferable, for reasons of economy, when many interpolated values are required.

To compute extrapolated values `nag_1d_cheb_interp` (e01aec), `nag_monotonic_interpolant` (e01bec) and `nag_1d_ratnl_interp` (e01rac) based on polynomials, piecewise polynomials and rational functions respectively, may be used. Extrapolation is not recommended in general, but can sometimes give acceptable results if it is to a point not far outside the data range, and only the few nearest data points are used in the process. `nag_1d_ratnl_interp` (e01rac) is most likely to be successful.

3.2.2 Interpolating function: data without derivatives

`nag_1d_cheb_interp` (e01aec) computes the Lagrange interpolating polynomial by a method (based on **Newton's formula with divided differences** (see Fröberg (1970)) which has proved numerically very stable. However, the likelihood of the polynomial having unwanted fluctuations, particularly near the ends of the data range when a moderate or large number of data points are used, should be remembered.

Such fluctuations of the polynomial can be avoided if you are at liberty to choose the x values at which to provide data points. In this case, a function from Chapter e02, namely `nag_1d_cheb_interp_fit` (e02afc), should be used in the manner and with the x values discussed in Section 3.2.2 in the e02 Chapter Introduction.

Usually however, when the whole of the data range is of interest, it is preferable to use a cubic spline as the interpolating function. `nag_1d_spline_interpolant` (e01bac) computes an interpolating cubic spline, using a particular choice for the set of knots which has proved generally satisfactory in practice. If you wish to choose a different set, a cubic spline function from Chapter e02, namely `nag_1d_spline_fit_knots` (e02bac), may be used in its interpolating mode, setting `spline.n = mdist + 4` and all elements of the argument `w` to unity.

The cubic spline does not always avoid unwanted fluctuations, especially when the data shows a steep slope close to a region of small slope, or when the data inadequately represents the underlying curve. In such cases, `nag_monotonic_interpolant` (e01bec) can be very useful. It derives a piecewise cubic polynomial (with first derivative continuity) which, between any adjacent pair of data points, either increases all the way, or decreases all the way (or stays constant). It is especially suited to data which is monotonic over the whole range.

In this function, the interpolating function is represented simply by its value and first derivative at the data points. Supporting functions compute its value and first derivative elsewhere, as well as its definite integral over an arbitrary interval. The other functions above provide the interpolating function either in Chebyshev-series form or in B-spline form (see Sections 2.2.1 and 2.2.2 in the e02 Chapter Introduction). Functions for evaluating, differentiating and integrating these forms are discussed in Section 3.7 in the e02 Chapter Introduction. The splines and other piecewise cubics will normally provide better estimates of the derivatives of the underlying function than will interpolating polynomials, at any rate away from the central part of the data range.

`nag_1d_ratnl_interp` (e01rac) computes an interpolating rational function. It is intended mainly for those cases where you know that this form of function is appropriate. However, it is also worth trying in cases where the other functions have proved unsatisfactory. `nag_1d_ratnl_eval` (e01rbc) is available to compute values of the function provided by `nag_1d_ratnl_interp` (e01rac).

3.2.3 Data containing derivatives

`nag_1d_cheb_interp` (e01aec) (see Section 3.2.2) can also compute the polynomial which, at each x_r , has not only a specified value y_r but also a specified value of each derivative up to order p_r .

3.3 Two Independent Variables

3.3.1 Data on a rectangular mesh

Given the value f_{qr} of the dependent variable f at the point (x_q, y_r) in the plane of the independent variables x and y , for each $q = 1, 2, \dots, m$ and $r = 1, 2, \dots, n$ (so that the points (x_q, y_r) lie at the $m \times n$ intersections of a rectangular mesh), `nag_2d_spline_interpolant` (e01dac) computes an interpolating bicubic spline, using a particular choice for each of the spline's knot-set. This choice, the same as in `nag_1d_spline_interpolant` (e01bac), has proved generally satisfactory in practice.

3.3.2 Arbitrary data

As remarked at the end of Section 2, special types of interpolating are required for this problem, which can often be difficult to solve satisfactorily. Two of the most successful are employed in `nag_2d_shep_interp` (e01sgc) and `nag_2d_triang_interp` (e01sjc) which, together with its evaluation function `nag_2d_shep_eval` (e01shc) and `nag_2d_triang_eval` (e01skc), is provided for the problem. Definitions can be found in the function documents. Both interpolants have first derivative continuity and are 'local', in that their value at any point depends only on data in the immediate neighbourhood of the point. This latter feature is necessary for large sets of data to avoid prohibitive computing time.

The relative merits of the two methods vary with the data and it is not possible to predict which will be the better in any particular case.


3.4 Three Independent Variables

3.4.1 Arbitrary data

The function `nag_3d_shep_interp` (e01tgc) and its evaluation function `nag_3d_shep_eval` (e01thc) are provided for interpolation of three-dimensional scattered data. As in the case of two independent variables, the method is local, and produces an interpolant with first derivative continuity. `nag_3d_shep_eval` (e01thc) allows evaluation of the interpolant and its first partial derivatives.

4 Decision Tree

Tree 1



5 Index

- Derivative,
of interpolant,
from `nag_monotonic_interpolant` (e01bec) `nag_monotonic_deriv` (e01bfc)
from `nag_2d_shep_interp` (e01sgc) `nag_2d_shep_eval` (e01shc)
from `nag_3d_shep_interp` (e01tgc) `nag_3d_shep_eval` (e01thc)
- Evaluation,
of interpolant,
from `nag_monotonic_interpolant` (e01bec) `nag_monotonic_evaluate` (e01bfc)
from `nag_1d_ratnl_interp` (e01rac) `nag_1d_ratnl_eval` (e01rbc)
from `nag_2d_shep_interp` (e01sgc) `nag_2d_shep_eval` (e01shc)
from `nag_3d_shep_interp` (e01tgc) `nag_3d_shep_eval` (e01thc)
from `nag_2d_triag_interp` (e01sjc) `nag_2d_triag_eval` (e01skc)
- Extrapolation,
one variable `nag_1d_cheb_interp` (e01aec)
one variable `nag_monotonic_interpolant` (e01bec)
one variable `nag_1d_ratnl_interp` (e01rac)

Integration (definite) of interpolant from nag_monotonic_interpolant (e01bec)	nag_monotonic_intg (e01bhc)
Interpolated values,	
one variable,	
from interpolant from nag_monotonic_interpolant (e01bec)	nag_monotonic_evaluate (e01bfc)
from interpolant from nag_monotonic_interpolant (e01bec) ...	nag_monotonic_deriv (e01bgc)
from rational function	nag_1d_ratnl_eval (e01rbc)
three variables,	
from interpolant from nag_3d_shep_interp (e01tgc)	nag_3d_shep_eval (e01thc)
two variables,	
from interpolant from nag_2d_shep_interp (e01sgc)	nag_2d_shep_eval (e01shc)
from interpolant from nag_2d_triang_interp (e01sjc)	nag_2d_triang_eval (e01skc)
Interpolating function,	
one variable,	
cubic spline	nag_1d_spline_interpolant (e01bac)
other piecewise polynomial	nag_monotonic_interpolant (e01bec)
polynomial,	
data with or without derivatives	nag_1d_cheb_interp (e01aec)
rational function	nag_1d_ratnl_interp (e01rac)
three variables,	
modified Shepard method	nag_3d_shep_interp (e01tgc)
two variables,	
bicubic spline	nag_2d_spline_interpolant (e01dac)
modified Shepard method	nag_2d_shep_interp (e01sgc)
other piecewise polynomial	nag_2d_triang_interp (e01sjc)

6 Functions Withdrawn or Scheduled for Withdrawal

Withdrawn Function	Mark of Withdrawal	Replacement Function(s)
nag_2d_scat_interpolant (e01sac)	10	nag_2d_shep_interp (e01sgc) or nag_2d_triang_interp (e01sjc)
nag_2d_scat_eval (e01sbc)	10	nag_2d_shep_eval (e01shc) or nag_2d_triang_eval (e01skc)
nag_2d_scat_free (e01szc)	10	No replacement document required

7 References

- Dahlquist G and Björck Å (1974) *Numerical Methods* Prentice–Hall
- Fröberg C E (1970) *Introduction to Numerical Analysis* Addison–Wesley