

# NAG Fortran Library Routine Document

## G02CDF

**Note:** before using this routine, please read the Users' Note for your implementation to check the interpretation of ***bold italicised*** terms and other implementation-dependent details.

### 1 Purpose

G02CDF performs a simple linear regression with no constant, with dependent variable  $y$  and independent variable  $x$ , omitting cases involving missing values.

### 2 Specification

```
SUBROUTINE G02CDF(N, X, Y, XMISS, YMISS, RESULT, IFAIL)
INTEGER          N, IFAIL
real           X(N), Y(N), XMISS, YMISS, RESULT(21)
```

### 3 Description

The routine fits a straight line of the form

$$y = bx$$

to those of the data points

$$(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$$

that do not include missing values, such that

$$y_i = bx_i + e_i$$

for those  $(x_i, y_i)$   $i = 1, 2, \dots, n$  ( $n \geq 2$ ) which do not include missing values.

The routine eliminates all pairs of observations  $(x_i, y_i)$  which contain a missing value for either  $x$  or  $y$ , and then calculates the regression coefficient,  $b$ , and various other statistical quantities by minimizing the sum of the  $e_i^2$  over those cases remaining in the calculations.

The input data consists of the  $n$  pairs of observations  $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$  on the independent variable  $x$  and the dependent variable  $y$ .

In addition two values,  $xm$  and  $ym$ , are given which are considered to represent missing observations for  $x$  and  $y$  respectively. (See Section 7).

Let  $w_i = 0$ , if the  $i$ th observation of either  $x$  or  $y$  is missing, i.e., if  $x_i = xm$  and/or  $y_i = ym$ ; and  $w_i = 1$  otherwise, for  $i = 1, 2, \dots, n$ .

The quantities calculated are:

(a) Means:

$$\bar{x} = \frac{\sum_{i=1}^n w_i x_i}{\sum_{i=1}^n w_i}; \quad \bar{y} = \frac{\sum_{i=1}^n w_i y_i}{\sum_{i=1}^n w_i}.$$

(b) Standard deviations:

$$s_x = \sqrt{\frac{\sum_{i=1}^n w_i (x_i - \bar{x})^2}{\sum_{i=1}^n w_i - 1}}; \quad s_y = \sqrt{\frac{\sum_{i=1}^n w_i (y_i - \bar{y})^2}{\sum_{i=1}^n w_i - 1}}.$$

(c) Pearson product-moment correlation coefficient:

$$r = \frac{\sum_{i=1}^n w_i (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n w_i (x_i - \bar{x})^2 \sum_{i=1}^n w_i (y_i - \bar{y})^2}}.$$

(d) The regression coefficient,  $b$ :

$$b = \frac{\sum_{i=1}^n w_i x_i y_i}{\sum_{i=1}^n w_i x_i^2}.$$

(e) The sum of squares attributable to the regression,  $SSR$ , the sum of squares of deviations about the regression,  $SSD$ , and the total sum of squares,  $SST$ :

$$SST = \sum_{i=1}^n w_i y_i^2; \quad SSD = \sum_{i=1}^n w_i (y_i - bx_i)^2; \quad SSR = SST - SSD.$$

(f) The degrees of freedom attributable to the regression,  $DFR$ , the degrees of freedom of deviations about the regression,  $DFD$ , and the total degrees of freedom,  $DFT$ :

$$DFT = \sum_{i=1}^n w_i; \quad DFD = \sum_{i=1}^n w_i - 1; \quad DFR = 1.$$

(g) The mean square attributable to the regression,  $MSR$ , and the mean square of deviations about the regression,  $MSD$ :

$$MSR = SSR/DFR; \quad MSD = SSD/DFD.$$

(h) The  $F$ -value for the analysis of variance:

$$F = MSR/MSD.$$

(i) The standard error of the regression coefficient:

$$se(b) = \sqrt{\frac{MSD}{\sum_{i=1}^n w_i x_i^2}}.$$

(j) The  $t$ -value for the regression coefficient:

$$t(b) = \frac{b}{se(b)}.$$

(k) The number of observations used in the calculations:

$$n_c = \sum_{i=1}^n w_i.$$

## 4 References

Draper N R and Smith H (1985) *Applied Regression Analysis* (2nd Edition) Wiley

## 5 Parameters

- |    |                                                                                                                   |              |
|----|-------------------------------------------------------------------------------------------------------------------|--------------|
| 1: | N – INTEGER                                                                                                       | <i>Input</i> |
|    | <i>On entry:</i> the number, $n$ , of pairs of observations.                                                      |              |
|    | <i>Constraint:</i> $N \geq 2$ .                                                                                   |              |
| 2: | X(N) – <i>real</i> array                                                                                          | <i>Input</i> |
|    | <i>On entry:</i> X( $i$ ) must contain $x_i$ , for $i = 1, 2, \dots, n$ .                                         |              |
| 3: | Y(N) – <i>real</i> array                                                                                          | <i>Input</i> |
|    | <i>On entry:</i> Y( $i$ ) must contain $y_i$ , for $i = 1, 2, \dots, n$ .                                         |              |
| 4: | XMISS – <i>real</i>                                                                                               | <i>Input</i> |
|    | <i>On entry:</i> the value $xm$ , which is to be taken as the missing value for the variable $x$ (see Section 7). |              |

- 5: YMISS – *real* *Input*  
*On entry:* the value  $ym$ , which is to be taken as the missing value for the variable  $y$  (see Section 7).
- 6: RESULT(21) – *real* array *Output*  
*On exit:* the following information:  
 RESULT(1)  $\bar{x}$ , the mean value of the independent variable,  $x$ ;  
 RESULT(2)  $\bar{y}$ , the mean value of the dependent variable,  $y$ ;  
 RESULT(3)  $s_x$ , the standard deviation of the independent variable,  $x$ ;  
 RESULT(4)  $s_y$ , the standard deviation of the dependent variable,  $y$ ;  
 RESULT(5)  $r$ , the Pearson product-moment correlation between the independent variable  $x$  and the dependent variable,  $y$ ;  
 RESULT(6)  $b$ , the regression coefficient;  
 RESULT(7) the value 0.0;  
 RESULT(8)  $se(b)$ , the standard error of the regression coefficient;  
 RESULT(9) the value 0.0;  
 RESULT(10)  $t(b)$ , the  $t$ -value for the regression coefficient;  
 RESULT(11) the value 0.0;  
 RESULT(12)  $SSR$ , the sum of squares attributable to the regression;  
 RESULT(13)  $DFR$ , the degrees of freedom attributable to the regression;  
 RESULT(14)  $MSR$ , the mean square attributable to the regression;  
 RESULT(15)  $F$ , the  $F$ -value for the analysis of variance;  
 RESULT(16)  $SSD$ , the sum of squares of deviations about the regression;  
 RESULT(17)  $DFD$ , the degrees of freedom of deviations about the regression;  
 RESULT(18)  $MSD$ , the mean square of deviations about the regression;  
 RESULT(19)  $SST$ , the total sum of squares  
 RESULT(20)  $DFT$ , the total degrees of freedom;  
 RESULT(21)  $n_c$ , the number of observations used in the calculations.
- 7: IFAIL – INTEGER *Input/Output*  
*On entry:* IFAIL must be set to 0,  $-1$  or 1. Users who are unfamiliar with this parameter should refer to Chapter P01 for details.  
*On exit:* IFAIL = 0 unless the routine detects an error (see Section 6).  
 For environments where it might be inappropriate to halt program execution when an error is detected, the value  $-1$  or 1 is recommended. If the output of error messages is undesirable, then the value 1 is recommended. Otherwise, for users not familiar with this parameter the recommended value is 0. **When the value  $-1$  or 1 is used it is essential to test the value of IFAIL on exit.**

## 6 Error Indicators and Warnings

If on entry IFAIL = 0 or  $-1$ , explanatory error messages are output on the current error message unit (as defined by X04AAF).

Errors or warnings detected by the routine:

IFAIL = 1

On entry,  $N < 2$ .

IFAIL = 2

After observations with missing values were omitted, fewer than two cases remained.

IFAIL = 3

After observations with missing values were omitted, all remaining values of at least one of the variables  $x$  and  $y$  were identical.

## 7 Accuracy

The routine does not use *additional precision* arithmetic for the accumulation of scalar products, so there may be a loss of significant figures for large  $n$ .

Users are warned of the need to exercise extreme care in their selection of missing values. The routine treats all values in the inclusive range  $(1 \pm \text{ACC}) \times xm_j$ , where  $xm_j$  is the missing value for variable  $j$  specified by the user, and ACC is a machine-dependent constant (see the Users' Note for your implementation) as missing values for variable  $j$ .

The user must therefore ensure that the missing value chosen for each variable is sufficiently different from all valid values for that variable so that none of the valid values fall within the range indicated above.

If, in calculating  $F$  or  $t(b)$  (see Section 3), the numbers involved are such that the result would be outside the range of numbers which can be stored by the machine, then the answer is set to the largest quantity which can be stored as a *real* variable, by means of a call to X02ALF.

## 8 Further Comments

The time taken by the routine depends on  $n$  and the number of missing observations.

The routine uses a two-pass algorithm.

## 9 Example

The following program reads in eight observations on each of two variables, and then performs a simple linear regression with no constant, with the first variable as the independent variable, and the second variable as the dependent variable, omitting cases involving missing values (0.0 for the first variable, 99.0 for the second). Finally the results are printed.

### 9.1 Program Text

**Note:** the listing of the example program presented below uses *bold italicised* terms to denote precision-dependent details. Please read the Users' Note for your implementation to check the interpretation of these terms. As explained in the Essential Introduction to this manual, the results produced may not be identical for all implementations.

```
*      G02CDF Example Program Text
*      Mark 14 Revised.  NAG Copyright 1989.
*      .. Parameters ..
      INTEGER          N
      PARAMETER       (N=8)
      INTEGER          NIN, NOUT
      PARAMETER       (NIN=5,NOUT=6)
*      .. Local Scalars ..
      real            XM, YM
      INTEGER          I, IFAIL
*      .. Local Arrays ..
      real            RESULT(21), X(N), Y(N)
*      .. External Subroutines ..
      EXTERNAL        G02CDF
*      .. Executable Statements ..
      WRITE (NOUT,*) 'G02CDF Example Program Results'
*      Skip heading in data file
      READ (NIN,*)
      READ (NIN,*) (X(I),Y(I),I=1,N)
      WRITE (NOUT,*)
      WRITE (NOUT,*) ' Case      Independent      Dependent'
      WRITE (NOUT,*) 'number      variable        variable'
      WRITE (NOUT,*)
      WRITE (NOUT,99999) (I,X(I),Y(I),I=1,N)
      WRITE (NOUT,*)
*
*      Set up missing values
*
      XM = 0.0e0
      YM = 99.0e0
      IFAIL = 1
```

```

*
  CALL G02CDF(N,X,Y,XM,YM,RESULT,IFAIL)
*
  IF (IFAIL.NE.0) THEN
    WRITE (NOUT,99998) 'Routine fails, IFAIL =', IFAIL
  ELSE
    WRITE (NOUT,99997)
+   'Mean of independent variable           = ', RESULT(1)
    WRITE (NOUT,99997)
+   'Mean of dependent variable             = ', RESULT(2)
    WRITE (NOUT,99997)
+   'Standard deviation of independent variable = ', RESULT(3)
    WRITE (NOUT,99997)
+   'Standard deviation of dependent variable = ', RESULT(4)
    WRITE (NOUT,99997)
+   'Correlation coefficient                 = ', RESULT(5)
    WRITE (NOUT,*)
    WRITE (NOUT,99997)
+   'Regression coefficient                  = ', RESULT(6)
    WRITE (NOUT,99997)
+   'Standard error of coefficient           = ', RESULT(8)
    WRITE (NOUT,99997)
+   't-value for coefficient                  = ', RESULT(10)
    WRITE (NOUT,*)
    WRITE (NOUT,*) 'Analysis of regression table :-'
    WRITE (NOUT,*)
    WRITE (NOUT,*)
+   '      Source          Sum of squares  D.F.    Mean square    F-val
+ue'
    WRITE (NOUT,*)
    WRITE (NOUT,99996) 'Due to regression', (RESULT(I),I=12,15)
    WRITE (NOUT,99996) 'About regression', (RESULT(I),I=16,18)
    WRITE (NOUT,99996) 'Total          ', (RESULT(I),I=19,20)
    WRITE (NOUT,*)
    WRITE (NOUT,99995) 'Number of cases used = ', RESULT(21)
  END IF
  STOP
*
99999 FORMAT (1X,I4,2F15.4)
99998 FORMAT (1X,A,I2)
99997 FORMAT (1X,A,F8.4)
99996 FORMAT (1X,A,F14.4,F8.0,2F14.4)
99995 FORMAT (1X,A,F3.0)
  END

```

## 9.2 Program Data

G02CDF Example Program Data

|      |      |
|------|------|
| 1.0  | 20.0 |
| 0.0  | 15.5 |
| 4.0  | 28.3 |
| 7.5  | 45.0 |
| 2.5  | 24.5 |
| 0.0  | 10.0 |
| 10.0 | 99.0 |
| 5.0  | 31.2 |

## 9.3 Program Results

G02CDF Example Program Results

| Case number | Independent variable | Dependent variable |
|-------------|----------------------|--------------------|
| 1           | 1.0000               | 20.0000            |
| 2           | 0.0000               | 15.5000            |
| 3           | 4.0000               | 28.3000            |
| 4           | 7.5000               | 45.0000            |
| 5           | 2.5000               | 24.5000            |
| 6           | 0.0000               | 10.0000            |

```

7      10.0000      99.0000
8      5.0000      31.2000

```

```

Mean of independent variable      = 4.0000
Mean of dependent variable        = 29.8000
Standard deviation of independent variable = 2.4749
Standard deviation of dependent variable = 9.4787
Correlation coefficient            = 0.9799

```

```

Regression coefficient             = 6.5833
Standard error of coefficient      = 0.8046
t-value for coefficient            = 8.1816

```

Analysis of regression table :-

| Source            | Sum of squares | D.F. | Mean square | F-value |
|-------------------|----------------|------|-------------|---------|
| Due to regression | 4528.9493      | 1.   | 4528.9493   | 66.9392 |
| About regression  | 270.6307       | 4.   | 67.6577     |         |
| Total             | 4799.5800      | 5.   |             |         |

Number of cases used = 5.

---