

# NAG Fortran Library Routine Document

## G07CAF

**Note:** before using this routine, please read the Users' Note for your implementation to check the interpretation of **bold italicised** terms and other implementation-dependent details.

### 1 Purpose

G07CAF computes a  $t$ -test statistic to test for a difference in means between two Normal populations, together with a confidence interval for the difference between the means.

### 2 Specification

```

SUBROUTINE G07CAF(TAIL, EQUAL, NX, NY, XMEAN, YMEAN, XSTD, YSTD, CLEVEL,
1              T, DF, PROB, DL, DU, IFAIL)
INTEGER       NX, NY, IFAIL
real        XMEAN, YMEAN, XSTD, YSTD, CLEVEL, T, DF, PROB, DL, DU
CHARACTER*1   TAIL, EQUAL

```

### 3 Description

Consider two independent samples, denoted by  $X$  and  $Y$ , of size  $n_x$  and  $n_y$  drawn from two Normal populations with means  $\mu_x$  and  $\mu_y$ , and variances  $\sigma_x^2$  and  $\sigma_y^2$  respectively. Denote the sample means by  $\bar{x}$  and  $\bar{y}$  and the sample variances by  $s_x^2$  and  $s_y^2$  respectively.

G07CAF calculates a test statistic and its significance level to test the null hypothesis  $H_0 : \mu_x = \mu_y$ , together with upper and lower confidence limits for  $\mu_x - \mu_y$ . The test used depends on whether or not the two population variances are assumed to be equal.

1. It is assumed that the two variances are equal, that is  $\sigma_x^2 = \sigma_y^2$ .

The test used is the two sample  $t$ -test. The test statistic  $t$  is defined by;

$$t_{obs} = \frac{\bar{x} - \bar{y}}{s \sqrt{(1/n_x) + (1/n_y)}}$$

where  $s^2 = ((n_x - 1)s_x^2 + (n_y - 1)s_y^2)/(n_x + n_y - 2)$  is the pooled variance of the two samples.

Under the null hypothesis  $H_0$  this test statistic has a  $t$ -distribution with  $(n_x + n_y - 2)$  degrees of freedom.

The test of  $H_0$  is carried out against one of three possible alternatives;

$H_1 : \mu_x \neq \mu_y$ ; the significance level,  $p = P(t \geq |t_{obs}|)$ , i.e., a two-tailed probability.

$H_1 : \mu_x > \mu_y$ ; the significance level,  $p = P(t \geq t_{obs})$ , i.e., an upper tail probability.

$H_1 : \mu_x < \mu_y$ ; the significance level,  $p = P(t \leq t_{obs})$ , i.e., a lower tail probability.

Upper and lower  $100(1 - \alpha)\%$  confidence limits for  $\mu_x - \mu_y$  are calculated as:

$$(\bar{x} - \bar{y}) \pm t_{1-\alpha/2} s \sqrt{(1/n_x) + (1/n_y)}.$$

where  $t_{1-\alpha/2}$  is the  $100(1 - \alpha/2)$  percentage point of the  $t$ -distribution with  $(n_x + n_y - 2)$  degrees of freedom.

2. It is not assumed that the two variances are equal.

If the population variances are not equal the usual two sample  $t$ -statistic no longer has a  $t$ -distribution and an approximate test is used.

This problem is often referred to as the Behrens–Fisher problem, see Kendall and Stuart (1969). The test used here is based on Satterthwaites procedure. To test the null hypothesis the test statistic  $t'$  is used where

$$t'_{obs} = \frac{\bar{x} - \bar{y}}{se(\bar{x} - \bar{y})}$$

$$\text{where } se(\bar{x} - \bar{y}) = \sqrt{\frac{s_x^2}{n_x} + \frac{s_y^2}{n_y}}$$

A  $t$ -distribution with  $f$  degrees of freedom is used to approximate the distribution of  $t'$  where

$$f = \frac{se(\bar{x} - \bar{y})^4}{\frac{s_x^2/n_x^2}{(n_x - 1)} + \frac{s_y^2/n_y^2}{(n_y - 1)}}$$

The test of  $H_0$  is carried out against one of the three alternative hypotheses described above, replacing  $t$  by  $t'$  and  $t_{obs}$  by  $t'_{obs}$ .

Upper and lower  $100(1 - \alpha)\%$  confidence limits for  $\mu_x - \mu_y$  are calculated as:

$$(\bar{x} - \bar{y}) \pm t_{1-\alpha/2} se(x - \bar{y}).$$

where  $t_{1-\alpha/2}$  is the  $100(1 - \alpha/2)$  percentage point of the  $t$ -distribution with  $f$  degrees of freedom.

## 4 References

Johnson M G and Kotz A (1969) *The Encyclopedia of Statistics 2* Griffin

Kendall M G and Stuart A (1969) *The Advanced Theory of Statistics (Volume 1)* (3rd Edition) Griffin

Snedecor G W and Cochran W G (1967) *Statistical Methods* Iowa State University Press

## 5 Parameters

- 1: TAIL – CHARACTER\*1 *Input*  
*On entry:* indicates which tail probability is to be calculated, and thus which alternative hypothesis is to be used.  
 If TAIL = 'T', the two tail probability, i.e.,  $H_1 : \mu_x \neq \mu_y$ .  
 If TAIL = 'U', the upper tail probability, i.e.,  $H_1 : \mu_x > \mu_y$ .  
 If TAIL = 'L', the lower tail probability, i.e.,  $H_1 : \mu_x < \mu_y$ .  
*Constraint:* TAIL = 'T', 'U' or 'L'.
- 2: EQUAL – CHARACTER\*1 *Input*  
*On entry:* indicates whether the population variances are assumed to be equal or not.  
 If EQUAL = 'E', the population variances are assumed to be equal, that is  $\sigma_x^2 = \sigma_y^2$ .  
 If EQUAL = 'U', the population variances are not assumed to be equal.  
*Constraint:* EQUAL = 'E' or 'U'.
- 3: NX – INTEGER *Input*  
*On entry:* the size of the  $X$  sample,  $n_x$ .  
*Constraint:* NX  $\geq$  2.

- 4: NY – INTEGER *Input*  
*On entry:* the size of the  $Y$  sample,  $n_y$ .  
*Constraint:*  $NY \geq 2$ .
- 5: XMEAN – *real* *Input*  
*On entry:* the mean of the  $X$  sample,  $\bar{x}$ .
- 6: YMEAN – *real* *Input*  
*On entry:* the mean of the  $Y$  sample,  $\bar{y}$ .
- 7: XSTD – *real* *Input*  
*On entry:* the standard deviation of the  $X$  sample,  $s_x$ .  
*Constraint:*  $XSTD > 0.0$ .
- 8: YSTD – *real* *Input*  
*On entry:* the standard deviation of the  $Y$  sample,  $s_y$ .  
*Constraint:*  $YSTD > 0.0$ .
- 9: CLEVEL – *real* *Input*  
*On entry:* the confidence level,  $1 - \alpha$ , for the specified tail. For example  $CLEVEL = 0.95$  will give a 95% confidence interval.  
*Constraint:*  $0.0 < CLEVEL < 1.0$ .
- 10: T – *real* *Output*  
*On exit:* contains the test statistic,  $t_{obs}$  or  $t'_{obs}$ .
- 11: DF – *real* *Output*  
*On exit:* contains the degrees of freedom for the test statistic.
- 12: PROB – *real* *Output*  
*On exit:* contains the significance level, that is the tail probability,  $p$ , as defined by TAIL.
- 13: DL – *real* *Output*  
*On exit:* contains the lower confidence limit for  $\mu_x - \mu_y$ .
- 14: DU – *real* *Output*  
*On exit:* contains the upper confidence limit for  $\mu_x - \mu_y$ .
- 15: IFAIL – INTEGER *Input/Output*  
*On entry:* IFAIL must be set to 0, -1 or 1. Users who are unfamiliar with this parameter should refer to Chapter P01 for details.  
*On exit:*  $IFAIL = 0$  unless the routine detects an error (see Section 6).  
For environments where it might be inappropriate to halt program execution when an error is detected, the value -1 or 1 is recommended. If the output of error messages is undesirable, then the value 1 is recommended. Otherwise, for users not familiar with this parameter the recommended value is 0. **When the value -1 or 1 is used it is essential to test the value of IFAIL on exit.**

## 6 Error Indicators and Warnings

If on entry  $IFAIL = 0$  or  $-1$ , explanatory error messages are output on the current error message unit (as defined by X04AAF).

Errors or warnings detected by the routine:

$IFAIL = 1$

On entry,  $TAIL \neq 'T', 'U'$  or  $'L'$ ,  
 or  $EQUAL \neq 'E'$  or  $'U'$ ,  
 or  $NX < 2$ ,  
 or  $NY < 2$ ,  
 or  $XSTD \leq 0.0$ ,  
 or  $YSTD \leq 0.0$ ,  
 or  $CLEVEL \leq 0.0$ ,  
 or  $CLEVEL \geq 1.0$ .

## 7 Accuracy

The computed probability and the confidence limits should be accurate to approximately 5 significant figures.

## 8 Further Comments

The sample means and standard deviations can be computed using G01AAF.

## 9 Example

The following example program reads the two sample sizes and the sample means and standard deviations for two independent samples. The data is taken from page 116, Snedecor and Cochran (1967) from a test to compare two methods of estimating the concentration of a chemical in a vat. A test of the equality of the means is carried out first assuming that the two population variances are equal and then making no assumption about the equality of the population variances.

### 9.1 Program Text

**Note:** the listing of the example program presented below uses *bold italicised* terms to denote precision-dependent details. Please read the Users' Note for your implementation to check the interpretation of these terms. As explained in the Essential Introduction to this manual, the results produced may not be identical for all implementations.

```
*      G07CAF Example Program Text
*      Mark 15 Release. NAG Copyright 1991.
*      .. Parameters ..
      INTEGER          NIN, NOUT
      PARAMETER       (NIN=5,NOUT=6)
*      .. Local Scalars ..
real               CLEVEL, DF, DL, DU, PROB, T, XMEAN, XSTD, YMEAN,
+                   YSTD
      INTEGER          IFAIL, NX, NY
*      .. External Subroutines ..
      EXTERNAL        G07CAF
*      .. Executable Statements ..
      WRITE (NOUT,*) 'G07CAF Example Program Results'
*      Skip heading in data file
      READ (NIN,*)
      READ (NIN,*) NX, NY
      READ (NIN,*) XMEAN, YMEAN, XSTD, YSTD
      READ (NIN,*) CLEVEL
      IFAIL = 0
*
      CALL G07CAF('Two', 'Equal', NX, NY, XMEAN, YMEAN, XSTD, YSTD, CLEVEL, T, DF,
+              PROB, DL, DU, IFAIL)
*
```

```

WRITE (NOUT,*)
WRITE (NOUT,*) 'Assuming population variances are equal.'
WRITE (NOUT,*)
WRITE (NOUT,99999) 't test statistic = ', T
WRITE (NOUT,99998) 'Degrees of freedom = ', DF
WRITE (NOUT,99997) 'Significance level = ', PROB
WRITE (NOUT,99999)
+ 'Lower confidence limit for difference in means = ', DL
WRITE (NOUT,99999)
+ 'Upper confidence limit for difference in means = ', DU
WRITE (NOUT,*)
IFAIL = 0
*
CALL G07CAF('Two','Unequal',NX,NY,XMEAN,YMEAN,XSTD,YSTD,CLEVEL,T,
+          DF,PROB,DL,DU,IFAIL)
*
WRITE (NOUT,*)
WRITE (NOUT,*) 'No assumptions about population variances .'
WRITE (NOUT,*)
WRITE (NOUT,99999) 't test statistic = ', T
WRITE (NOUT,99997) 'Degrees of freedom = ', DF
WRITE (NOUT,99997) 'Significance level = ', PROB
WRITE (NOUT,99999)
+ 'Lower confidence limit for difference in means = ', DL
WRITE (NOUT,99999)
+ 'Upper confidence limit for difference in means = ', DU
STOP
*
99999 FORMAT (1X,A,F10.4)
99998 FORMAT (1X,A,F8.1)
99997 FORMAT (1X,A,F8.4)
END

```

## 9.2 Program Data

G07CAF Example Program Data

```

4 8
25.0 21.0
0.8185 4.2083
0.95

```

## 9.3 Program Results

G07CAF Example Program Results

Assuming population variances are equal.

```

t test statistic =      1.8403
Degrees of freedom =      10.0
Significance level =      0.0955
Lower confidence limit for difference in means =      -0.8429
Upper confidence limit for difference in means =      8.8429

```

No assumptions about population variances .

```

t test statistic =      2.5922
Degrees of freedom =      7.9925
Significance level =      0.0320
Lower confidence limit for difference in means =      0.4410
Upper confidence limit for difference in means =      7.5590

```

---