

# NAG Fortran Library Routine Document

## D01ARF

**Note:** before using this routine, please read the Users' Note for your implementation to check the interpretation of *bold italicised* terms and other implementation-dependent details.

### 1 Purpose

D01ARF computes definite and indefinite integrals over a finite range to a specified relative or absolute accuracy, using the method described by Patterson (1968b).

### 2 Specification

```

SUBROUTINE D01ARF(A, B, FUN, RELACC, ABSACC, MAXRUL, IPARM, ACC, ANS, N,
1              ALPHA, IFAIL)
  INTEGER      MAXRUL, IPARM, N, IFAIL
  real       A, B, FUN, RELACC, ABSACC, ACC, ANS, ALPHA(390)
  EXTERNAL    FUN

```

### 3 Description

This routine evaluates definite and indefinite integrals of the form:

$$\int_a^b f(t) dt$$

using the method described by Patterson (1968b).

#### 3.1 Definite Integrals

In this case the routine must be called with  $IPARM = 0$ . By linear transformation the integral is changed to

$$I = \int_{-1}^{+1} F(x) dx$$

where

$$F(x) = \frac{b-a}{2} f\left(\frac{b+a+(b-a)x}{2}\right)$$

and is then approximated by an  $n$ -point quadrature rule

$$I = \sum_{k=1}^n w_k F(x_k)$$

where  $w_k$  are the weights and  $x_k$  are the abscissae.

The routine uses a family of 9 interlacing rules based on the optimal extension of the three-point Gauss rule. These rules use 1, 3, 7, 15, 31, 63, 127, 255 and 511 points and have respective polynomial integrating degrees 1, 5, 11, 23, 47, 95, 191, 383 and 767. Each rule has the property that the next in sequence includes all the points of its predecessor and has the greatest possible increase in integrating degree.

The integration method is based on the successive application of these rules until the absolute value of the difference of two successive results differs by not more than  $ABSACC$ , or relatively by not more than  $RELACC$ . The result of the last rule used is taken as the value of the integral ( $ANS$ ), and the absolute difference of the results of the last two rules used is taken as an estimate of the absolute error ( $ACC$ ). Due to their interlacing form no integrand evaluations are wasted in passing from one rule to the next.

### 3.2 Indefinite Integrals

Suppose the value of the integral

$$\int_c^d f(t) dt$$

is required for a number of sub-intervals  $[c, d]$ , all of which lie in a interval  $[a, b]$ .

In this case the routine should first be called with the parameter  $IPARM = 1$  and the interval set to  $[a, b]$ . The routine then calculates the integral over  $[a, b]$  **and** the Legendre expansion of the integrand, using the same integrand values. If the routine is subsequently called with  $IPARM = 2$  and the interval set to  $[c, d]$ , the integral over  $[c, d]$  is calculated by analytical integration of the Legendre expansion, without further evaluations of the integrand.

For the interval  $[-1, 1]$  the expansion takes the form

$$F(x) = \sum_{i=0}^{\infty} \alpha_i P_i(x)$$

where  $P_i(x)$  is the order  $i$  Legendre polynomial. Assuming that the integral over the full range  $[-1, 1]$  was evaluated to the required accuracy using an  $n$ -point rule, then the coefficients

$$\alpha_i = \frac{1}{2}(2i - 1) \int_{-1}^{+1} P_i(x) F(x) dx, \quad i = 0, 1, \dots, m$$

are evaluated by that same rule, up to

$$m = (3n - 1)/4.$$

The accuracy for indefinite integration should be of the same order as that obtained for the definite integral over the full range. The indefinite integrals will be exact when  $F(x)$  is a polynomial of degree  $\leq m$ .

## 4 References

Patterson T N L (1968b) The Optimum addition of points to quadrature formulae *Math. Comput.* **22** 847–856

## 5 Parameters

- 1: A – *real* *Input*  
*On entry:* the lower limit of integration,  $a$ .
- 2: B – *real* *Input*  
*On entry:* the upper limit of integration,  $b$ . It is not necessary that  $a < b$ .
- 3: FUN – *real* FUNCTION, supplied by the user. *External Procedure*  
 FUN must evaluate the integrand  $f$  at a specified point.  
 Its specification is:

<pre> <b>real</b> FUNCTION FUN(X) <b>real</b>           X           </pre>
1: X – <i>real</i> <span style="float: right;"><i>Input</i></span> <i>On entry:</i> the point in $[a, b]$ at which the integrand must be evaluated.

FUN must be declared as EXTERNAL in the (sub)program from which D01ARF is called. Parameters denoted as *Input* must **not** be changed by this procedure.

If  $IPARM = 2$ , FUN is not called.

- 4: RELACC – *real* *Input*  
*On entry:* the relative accuracy required. If convergence according to absolute accuracy is required, RELACC should be set to zero (but see also Section 7). If RELACC < 0.0, its absolute value is used.  
 If IPARM = 2, RELACC is not used.
- 5: ABSACC – *real* *Input*  
*On entry:* the absolute accuracy required. If convergence according to relative accuracy is required, ABSACC should be set to zero (but see also Section 7). If ABSACC < 0.0, its absolute value is used.  
 If IPARM = 2, ABSACC is not used.
- 6: MAXRUL – INTEGER *Input*  
*On entry:* the maximum number of successive rules that may be used.  
*Constraint:*  $1 \leq \text{MAXRUL} \leq 9$ . If MAXRUL is outside these limits, the value 9 is assumed.  
 If IPARM = 2, MAXRUL is not used..
- 7: IPARM – INTEGER *Input*  
*On entry:* IPARM indicates the task to be performed by the routine:  
     if IPARM = 0, only the definite integral over  $[a, b]$  is evaluated.  
     if IPARM = 1, as well as the definite integral, the expansion of the integrand in Legendre polynomials over  $[a, b]$  is calculated, using the same values of the integrand as used to compute the integral. The expansion coefficients, and some other quantities, are returned in ALPHA for later use in computing indefinite integrals.  
     if IPARM = 2,  $f(t)$  is integrated analytically over  $[a, b]$  using the previously computed expansion, stored in ALPHA. No further evaluations of the integrand are required. The routine must previously have been called with IPARM = 1 and the interval  $[a, b]$  must lie within that specified for the previous call. In this case only the arguments A, B, IPARM, ANS, ALPHA and IFAIL are used.  
*Constraint:* IPARM = 0, 1 or 2.
- 8: ACC – *real* *Output*  
*On exit:* if IPARM = 0 or 1, ACC contains the absolute value of the difference between the last two successive estimates of the integral. This may be used as a measure of the accuracy actually achieved.  
 If IPARM = 2, ACC is not used.
- 9: ANS – *real* *Output*  
*On exit:* the estimated value of the integral.
- 10: N – INTEGER *Output*  
*On exit:* when IPARM = 0 or 1, N contains the number of integrand evaluations used in the calculation of the integral.  
 If IPARM = 2, N is not used.
- 11: ALPHA(390) – *real* array *Input/Output*  
*On entry:* if IPARM = 2, ALPHA must contain the coefficients of the Legendre expansions of the integrand, as returned by a previous call of D01ARF with IPARM = 1 and a range containing the present range. If IPARM = 0 or 1, ALPHA need not be set on entry.

*On exit:* if  $IPARM = 1$ , the first  $m$  elements of ALPHA hold the coefficients of the Legendre expansion of the integrand, and the value of  $m$  is stored in ALPHA(390). ALPHA must not be changed between a call with  $IPARM = 1$  and subsequent calls with  $IPARM = 2$ .

If  $IPARM = 2$ , the first  $m$  elements of ALPHA are unchanged on exit.

12: IFAIL – INTEGER

*Input/Output*

*On entry:* IFAIL must be set to 0, -1 or 1. Users who are unfamiliar with this parameter should refer to Chapter P01 for details.

*On exit:* IFAIL = 0 unless the routine detects an error (see Section 6).

For environments where it might be inappropriate to halt program execution when an error is detected, the value -1 or 1 is recommended. If the output of error messages is undesirable, then the value 1 is recommended. Otherwise, because for this routine the values of the output parameters may be useful even if  $IFAIL \neq 0$  on exit, the recommended value is -1. **When the value -1 or 1 is used it is essential to test the value of IFAIL on exit.**

## 6 Error Indicators and Warnings

If on entry  $IFAIL = 0$  or  $-1$ , explanatory error messages are output on the current error message unit (as defined by X04AAF).

Errors or warnings detected by the routine:

IFAIL = 1

If  $IPARM = 0$  or  $1$ , this indicates that all MAXRUL rules have been used and the integral has not converged to the accuracy requested. In this case ANS contains the last approximation to the integral, and ACC contains the difference between the last two approximations. To check this estimate of the integral, D01ARF could be called again to evaluate

$$\int_a^b f(t) dt \quad \text{as} \quad \int_a^c f(t) dt + \int_c^b f(t) dt \quad \text{for some } a < c < b.$$

If  $IPARM = 2$ , this indicates failure of convergence during the run with  $IPARM = 1$  in which the Legendre expansion was created.

IFAIL = 2

On entry,  $IPARM < 0$  or  $IPARM > 2$ .

IFAIL = 3

The routine is called with  $IPARM = 2$  but a previous call with  $IPARM = 1$  has been omitted or was invoked with an integration interval of length zero.

IFAIL = 4

On entry, with  $IPARM = 2$ , the interval for indefinite integration is not contained within the interval specified when the routine was previously called with  $IPARM = 1$ .

## 7 Accuracy

The relative or absolute accuracy required is specified by the user in the variables RELACC or ABSACC. The routine will terminate whenever either the relative accuracy specified by RELACC or the absolute accuracy specified by ABSACC is reached. One or other of these criteria may be 'forced' by setting the parameter for the other to zero. If both RELACC and ABSACC are specified as zero, then the routine uses the value  $10.0 \times (\textit{machine precision})$  for RELACC.

If on exit  $IFAIL = 0$ , then it is likely that the result is correct to one or other of these accuracies. If on exit  $IFAIL = 1$ , then it is likely that neither of the requested accuracies has been reached.

When the user has no prior idea of the magnitude of the integral, it is possible that an unreasonable accuracy may be requested, e.g., a relative accuracy for an integral which turns out to be zero, or a small absolute accuracy for an integral which turns out to be very large. Even if failure is reported in such a case, the value of the integral may still be satisfactory. The device of setting the other ‘unused’ accuracy parameter to a small positive value (e.g.,  $10^{-9}$  for an implementation of 11-digit precision) rather than zero, may prevent excessive calculation in such a situation.

To avoid spurious convergence, it is recommended that relative accuracies larger than about  $10^{-3}$  be avoided.

## 8 Further Comments

The time taken by the routine depends on the complexity of the integrand and the accuracy required.

This routine uses the Patterson method over the whole integration interval and should therefore be suitable for well behaved functions. However, for very irregular functions it would be more efficient to submit the differently behaved regions separately for integration.

## 9 Example

The program evaluates the following integrals:

- (i) Definite integral only (IPARM = 0) for

$$\int_0^1 \frac{4}{1+x^2} dx \quad (\text{ABSACC} = 10^{-5}).$$

- (ii) Definite integral together with expansion coefficients (IPARM = 1) for

$$\int_1^2 \sqrt[8]{x} dx \quad (\text{ABSACC} = 10^{-5}).$$

- (iii) Indefinite integral using previous expansion (IPARM = 2) for

$$\int_{1.2}^{1.8} \sqrt[8]{x} dx \quad (\text{ABSACC} = 10^{-5}).$$

### 9.1 Program Text

**Note:** the listing of the example program presented below uses *bold italicised* terms to denote precision-dependent details. Please read the Users’ Note for your implementation to check the interpretation of these terms. As explained in the Essential Introduction to this manual, the results produced may not be identical for all implementations.

```
*      D01ARF Example Program Text
*      Mark 16 Revised. NAG Copyright 1993.
*      .. Parameters ..
INTEGER          NOUT
PARAMETER       (NOUT=6)
INTEGER          MAXRUL
PARAMETER       (MAXRUL=0)
*      .. Local Scalars ..
real           A, ABSACC, ACC, ANS, B, RELACC
INTEGER          IFAIL, IPARM, N
*      .. Local Arrays ..
real           ALPHA(390)
*      .. External Functions ..
real           F1, F2
EXTERNAL        F1, F2
*      .. External Subroutines ..
EXTERNAL        D01ARF
*      .. Executable Statements ..
WRITE (NOUT,*) 'D01ARF Example Program Results'
RELACC = 0.0e0
ABSACC = 1.0e-5
*      Definite integral of F1(x) - no expansion
IPARM = 0
```

```

IFAIL = 1
A = 0.0e0
B = 1.0e0
WRITE (NOUT,*)
WRITE (NOUT,*) 'Definite integral of 4/(1+x*x) over (0,1)'
*
CALL D01ARF(A,B,F1,RELACC,ABSACC,MAXRUL,IPARM,ACC,ANS,N,ALPHA,
+          IFAIL)
*
IF (IFAIL.NE.0) WRITE (NOUT,99997) 'D01ARF fails. IFAIL =', IFAIL
IF (IFAIL.LE.1) THEN
  WRITE (NOUT,99999) 'Estimated value of the integral =', ANS
  WRITE (NOUT,99998) 'Estimated absolute error =', ACC
  WRITE (NOUT,99997) 'Number of points used =', N
END IF
*
Definite integral of F2(x) - with expansion
IPARM = 1
IFAIL = 1
A = 1.0e0
B = 2.0e0
WRITE (NOUT,*)
WRITE (NOUT,*) 'Definite integral of x**(1/8) over (1,2)'
*
CALL D01ARF(A,B,F2,RELACC,ABSACC,MAXRUL,IPARM,ACC,ANS,N,ALPHA,
+          IFAIL)
*
IF (IFAIL.NE.0) WRITE (NOUT,99997) 'D01ARF fails. IFAIL =', IFAIL
IF (IFAIL.LE.1) THEN
  WRITE (NOUT,99999) 'Estimated value of the integral =', ANS
  WRITE (NOUT,99998) 'Estimated absolute error =', ACC
  WRITE (NOUT,99997) 'Number of points used =', N
END IF
*
Indefinite integral of F2(x)
IPARM = 2
IFAIL = 0
A = 1.2e0
B = 1.8e0
WRITE (NOUT,*)
WRITE (NOUT,*) 'Indefinite integral of x**(1/8) over (1.2,1.8)'
*
CALL D01ARF(A,B,F2,RELACC,ABSACC,MAXRUL,IPARM,ACC,ANS,N,ALPHA,
+          IFAIL)
*
WRITE (NOUT,99999) 'Estimated value of the integral =', ANS
STOP
*
99999 FORMAT (1X,A,F9.5)
99998 FORMAT (1X,A,e10.2)
99997 FORMAT (1X,A,I4)
END
*
real FUNCTION F1(X)
*
.. Scalar Arguments ..
real X
*
.. Executable Statements ..
F1 = 4.0e0/(1.0e0+X*X)
RETURN
END
*
real FUNCTION F2(X)
*
.. Scalar Arguments ..
real X
*
.. Executable Statements ..
F2 = X**0.125e0
RETURN
END

```

## 9.2 Program Data

None.

### 9.3 Program Results

D01ARF Example Program Results

Definite integral of  $4/(1+x*x)$  over (0,1)  
Estimated value of the integral = 3.14159  
Estimated absolute error = 0.18E-07  
Number of points used = 15

Definite integral of  $x^{1/8}$  over (1,2)  
Estimated value of the integral = 1.04979  
Estimated absolute error = 0.59E-06  
Number of points used = 7

Indefinite integral of  $x^{1/8}$  over (1.2,1.8)  
Estimated value of the integral = 0.63073

---