

## Moment Singularity Analysis of Vibration Spectra\*†

MELVIN LAX AND JOEL L. LEBOWITZ

*Department of Physics, Syracuse University, Syracuse, New York*

(Received July 7, 1954)

The frequency distribution of a crystal is approximated by combining Van Hove's determination of its analytical nature and Montroll's method of moments. The function  $G(\omega^2)$  is represented by an expression with the correct behavior at the singularities and at the maximum and minimum frequencies. The behavior between singular points is adjusted smoothly by leaving  $n$  undetermined parameters. These parameters are then fixed by using the correct first  $n$  moments. As a test, this procedure was applied to the two-dimensional square lattice with nearest and next nearest neighbor interactions, solved exactly for a particular case by Montroll. The approximated distribution function had the right form at the end points, contained terms of the appropriate logarithmic form, and a jump function (with known coefficients). It also included Legendre polynomials with unknown coefficients, which were determined by moments. The difference between the exact and approximate distribution functions was a few percent using only the zeroth moment (normalization). Using higher moments produced a gradual increase in accuracy.

## I. INTRODUCTION

THE normal modes of a lattice are eigenvectors of a matrix  $M_{ij}^{\alpha,\beta}$  denoted simply by  $\mathbf{M}$ , where  $i, j$  are indices of the cell and  $\alpha, \beta$  are indices of particles in a cell. Using the translational symmetry of the lattice, the problem is customarily reduced to the diagonalization of a matrix  $M^{\alpha,\beta}(\mathbf{k})$ , denoted briefly by  $\mathbf{M}(\mathbf{k})$ , where  $\mathbf{k}$  is the propagation constant of the wave, and the index  $\alpha$  (or  $\beta$ ) takes on  $lZ$  different values, where  $l$  is the dimension of the space and  $Z$  the number of atoms per cell. The elements of  $\mathbf{M}(\mathbf{k})$  are periodic functions of  $\mathbf{k}$ , with the periodicity of the reciprocal lattice.

The normalized density of eigenvalues  $\omega^2$  associated with the matrix  $\mathbf{M}$  can be defined by

$$G(\omega^2) = \langle \delta(\omega^2 - \mathbf{M}) \rangle, \quad (1)$$

where the average of a matrix  $\langle \rangle$  is the trace of that matrix divided by its dimensionality. Using the uniform spacing of the propagation constant over one Brillouin zone, Eq. (1) can be reduced to

$$G(\omega^2) = \int \langle \delta[\omega^2 - \mathbf{M}(\mathbf{k})] \rangle d\mathbf{k} / \int d\mathbf{k}, \quad (2)$$

where

$$\begin{aligned} \langle \delta[\omega^2 - \mathbf{M}(\mathbf{k})] \rangle &= (lZ)^{-1} \text{trace} \{ \delta[\omega^2 - \mathbf{M}(\mathbf{k})] \} \\ &= (lZ)^{-1} \sum_{n=1}^{lZ} \delta[\omega^2 - \omega_n^2(\mathbf{k})]. \end{aligned}$$

The relation between the frequency distribution  $g(\omega)$  and the distribution function of the frequency squared  $G(\omega^2)$  is

$$g(\omega) = 2\omega G(\omega^2). \quad (3)$$

\* A summary of the essential results of this paper was presented to the Washington meeting of the American Physical Society, April, 1954.

† Supported in part by the Office of Naval Research.

Since the integral in Eq. (2) cannot be evaluated explicitly except for a few special cases,<sup>1-3</sup> many approximate methods<sup>4</sup> have been used to study  $g(\omega)$ . Blackman<sup>5</sup> approximates  $g(\omega)$  by calculating the frequencies at a large number of points in the Brillouin zone and finding their distribution. This requires a great deal of labor, which has to be repeated for every set of force constants. Houston<sup>6</sup> finds the distribution along special lines in reciprocal space and interpolates for the rest of the Brillouin zone. This method introduces some spurious singularities in  $g(\omega)$ .<sup>7</sup> Montroll uses the moments of  $g(\omega)$  to determine the coefficients of the initial terms in a power series expansion of  $g(\omega)$ . We shall discuss Montroll's<sup>8,9</sup> moment method in some detail as it forms part of the basis of our work.<sup>10</sup>

From Eq. (1), the  $n$ th moment of  $G(x)$  is given by

$$m_n = \int_0^1 x^n G(x) dx = \langle \mathbf{M}^n \rangle, \quad (4)$$

or, taking explicitly the average over the propagation constant<sup>11</sup> as in Eq. (2), by

$$m_n = \int \langle \mathbf{M}^n(\mathbf{k}) \rangle d\mathbf{k} / \int d\mathbf{k}. \quad (5)$$

<sup>1</sup> E. W. Montroll, J. Chem. Phys. **15**, 575 (1947).

<sup>2</sup> W. A. Bowers and H. B. Rosenstock, J. Chem. Phys. **18**, 1056 (1950).

<sup>3</sup> G. F. Newell, J. Chem. Phys. **21**, 1877 (1953); H. B. Rosenstock, J. Chem. Phys. **21**, 2064 (1953); H. B. Rosenstock and G. F. Newell **21**, 1607 (1953); H. B. Rosenstock and H. M. Rosenstock **21**, 1608 (1953).

<sup>4</sup> A discussion of some of the methods together with references is given by A. C. Menzies, Repts. Progr. Phys. **16**, 83 (1953).

<sup>5</sup> M. Blackman, Repts. Progr. Phys. **8**, 11 (1941).

<sup>6</sup> W. V. Houston, Revs. Modern Phys. **20**, 161 (1948).

<sup>7</sup> T. Nakamura, Prog. Theoret. Phys. **5**, 213 (1950).

<sup>8</sup> E. W. Montroll, J. Chem. Phys. **10**, 218 (1942).

<sup>9</sup> E. W. Montroll, J. Chem. Phys. **11**, 481 (1943).

<sup>10</sup> For convenience we use units in which the maximum frequency of vibrations  $\omega_L$  is unity so that  $x = \omega^2 = (\omega/\omega_L)^2$  covers the range  $0 \leq x \leq 1$ .

<sup>11</sup> T. H. Walnut, J. Chem. Phys. **22**, 692 (1954), calculates the moments directly from  $\mathbf{M}$ .

Montroll was the first to recognize that the moments  $m_n$  are easily computed from traces of powers of  $\mathbf{M}(\mathbf{k})$  via Eq. (5). He showed that the first  $N$  moments can be used to construct the best approximate distribution function in the sense of least squares, for approximating functions expressed in a power series or a polynomial series containing  $N$  terms.<sup>12</sup> For these two choices of expansion, the least squares criterion is equivalent to the requirement that the first  $N$  moments of the approximating and exact distribution agree.<sup>13</sup> The rate of convergence of such an expansion is determined by the smoothness of the exact function. Hence, if  $G(x)$  is smooth, the use of even a few moments could be expected to give a good approximation.

However, it was found by Montroll,<sup>1</sup> when he did an exact calculation for a two-dimensional square lattice with nearest and next nearest neighbor interactions, that the frequency distribution function was not smooth but contained two logarithmic singularities. Smollett<sup>14</sup> extended Montroll's result to the case of a two-dimensional ionic lattice, taking into account the long-range Coulomb forces between ions. Similar results were obtained by Bowers and Rosenstock<sup>2</sup> for the frequency distribution of vibrations perpendicular to the plane of the lattice. Van Hove<sup>15</sup> subsequently showed that these singularities, far from being accidental, are a necessary consequence of the periodic structure of the lattice. They arise from those vibrations in the neighborhood of critical points  $\mathbf{k}_c$  for which  $|\text{grad}x_n(\mathbf{k}_c)|=0$ . The singularities in  $G(x)$  occur at  $x_c=x_n(\mathbf{k}_c)$ , where  $x_n(\mathbf{k})=\omega_n^2(\mathbf{k})$ . In two dimensions  $G(x)$  will have at least one logarithmic singularity. In three dimensions there will be at least three critical points  $x_c$ , where while  $G(x)$  remains continuous, its derivative  $G'(x_c)$  has an inverse square root singularity.

These considerations show that a smooth approximating function  $G_a(x)$ , such as a linear combination of polynomials which has the correct first  $N$  moments, would converge slowly to  $G(x)$  with increasing  $N$ . Hence a large number of moments, and a corresponding large amount of labor, would be needed to get a good fit to  $G(x)$ .

We decided, therefore, to take explicitly into account the analytic nature of  $G(x)$  in constructing  $G_a(x)$ .  $G_a(x)$  is represented by an expression with the correct behavior at the singularities and at the end points. The behavior between singular points is adjusted smoothly by leaving  $n$  undetermined parameters to be

fitted by the moments.<sup>16</sup> In this way the moments are used for approximating a smooth function so that good agreement to  $G(x)$  might be expected, even when, only a small number of moments are used. Essentially the same procedure was suggested independently by Rosenstock,<sup>17</sup> who applied it to the body-centered and face-centered cubic lattices. See Sec. III for further discussion of Rosenstock's results.

## II. TWO-DIMENSIONAL SQUARE LATTICE

As a test, this procedure, which we call the moment-singularity method was applied to a two-dimensional monatomic square lattice with neighbor and next neighbor interactions. The distribution function obtained by this procedure will be compared (a) with the exact distribution calculated by Montroll,<sup>1</sup> (b) with the distribution obtained by the unmodified method of moments.

For this lattice,  $\mathbf{M}(\mathbf{k})$  is a two-by-two matrix:<sup>18</sup>

$$\mathbf{M}(\mathbf{k}) = \frac{1}{2} \begin{pmatrix} 1 + (t-1)c_1 - tc_1c_2 & ts_1s_2 \\ ts_1s_2 & 1 + (t-1)c_2 - tc_1c_2 \end{pmatrix}, \quad (6)$$

where  $c_i = \cos k_i$ ,  $s_i = \sin k_i$ ,  $i=1,2$ ,  $t = (1 + \alpha/2\gamma)^{-1}$ , and  $\alpha$  and  $\gamma$  are neighbor and next neighbor forces, respectively. The two roots of the secular equation are

$$x_{\pm} = \frac{1}{4} [2t(1 - c_1c_2) + (1-t)(2 - c_1 - c_2) \pm \frac{1}{4} [4t^2s_1^2s_2^2 + (1-t)^2(c_1 - c_2)^2]]^{\frac{1}{2}}. \quad (7)$$

The  $n$ th moment of  $G(x)$  can be found from Eq. (5) by integrating the trace  $M^n(k)$  over the Brillouin zone,  $-\pi < k_i, k_j \leq \pi$ , and dividing the result by  $2(2\pi)^2$ . Thus, the first moment  $m_1$ , is

$$m_1 = \int_{-\pi}^{\pi} dk_1 \int_{-\pi}^{\pi} dk_2 (16\pi^2)^{-1} \times [2 - (1-t)(c_1 + c_2) - 2tc_1c_2] = \frac{1}{2}. \quad (8)$$

Montroll has calculated the first six even moments  $\mu_{2n}$  of  $g(\omega)$ , in a slightly different way. These are related to the  $m_n$  by

$$\mu_{2n} = \int_0^1 \omega^{2n} g(\omega) d\omega = \int_0^1 x^n G(x) dx = m_n, \quad (9)$$

and their values are

$$\begin{aligned} m_0 &= 1, & m_1 &= 1/2, & m_2 &= (3 - 2t + 2t^2)/8; \\ m_3 &= (5 - 6t + 6t^2)/16; \\ m_4 &= (35 - 60t + 77t^2 - 34t^3 + 17t^4)/128; \\ m_5 &= (462 - 980t + 960t^2 - 265t^3 + 80t^4 - 5t^5)/1024. \end{aligned} \quad (10)$$

<sup>16</sup> The fitting between singular points may also be done by any other method, e.g., Blackman's which yields sufficiently accurate results away from the singularities.

<sup>17</sup> H. B. Rosenstock, Phys. Rev. **95**, 617 (1954). A paper reporting this work is in preparation.

<sup>18</sup> For comparison with notation in reference 1:  $k_i \rightarrow \Phi_i$ ,  $x \rightarrow f^2$ ,  $G(x) \rightarrow (2f)^{-1} \omega_L G(f)$ , we measure the frequency in such units that the largest frequency  $\omega_L = 1$ .

<sup>12</sup> The use of polynomials as expansion functions will always lead to the same result as the use of a power series, for the same number of terms. The use of orthogonal polynomials greatly simplifies the arithmetic.

<sup>13</sup> If an expansion is made in functions other than polynomials, e.g., an  $N$ -term Fourier series, the method of equating moments and the least squares method are distinct for finite  $N$ . While knowledge of the first  $N$  moments determines a Fourier expansion to  $N$  terms, if the moment conditions are applied, this knowledge is insufficient to lead to a least squares solution.

<sup>14</sup> M. Smollett, Proc. Phys. Soc. (London) **A65**, 109 (1952).

<sup>15</sup> L. Van Hove, Phys. Rev. **89**, 1189 (1953).

TABLE I. Value of the coefficients of  $A_k^{(N)}$  of the Legendre polynomials in the expansion of the nonsingular part of distribution function.

$N \backslash k$	$A_0^{(N)}$	$A_1^{(N)}$	$A_2^{(N)}$	$A_3^{(N)}$	$A_4^{(N)}$	$A_5^{(N)}$	$A_6^{(N)}$	$A_7^{(N)}$
$G_a^{(1)}$	-0.1306	-0.1599	0.1568					
$G_a^{(2)}$	-0.1306	-0.2620	0.1568	0.1021				
$G_a^{(3)}$	-0.1306	-0.2620	0.1409	0.1021	0.0159			
$G_a^{(4)}$	-0.1306	-0.2620	0.1409	0.1388	0.0159	-0.0368		
$G_a^{(5)}$	-0.1306	-0.2620	0.1409	0.1388	0.0536	-0.0368	-0.0378	
$G_a^{(6)}$	-0.1306	-0.2620	0.1409	0.1388	0.0536	-0.0429	-0.0378	0.0061

An explicit form for all higher moments was found by Walnut.<sup>11</sup>

Because of the symmetry of the problem it is sufficient to work in one quadrant of the Brillouin zone,  $0 \leq k_1, k_2 \leq \pi$ . In this quadrant the critical points of  $x_{\pm}$  are at

$$(k_1, k_2) = (0, 0), (0, \pi), (\pi, 0), (\pi, \pi), \text{ when } 0 < t < 1/5. \quad (11)$$

The values of  $x$ , in the two branches  $x_{\pm}(\mathbf{k})$ , at these critical points are  $(0, 1, 1, 1-t)$  and  $(0, t, t, 1-t)$ , respectively. For  $t$  greater than  $1/5$  but less than  $1/2$ ,<sup>19</sup> the  $x_+$  branch has one additional critical point at  $\{\cos^{-1}[(t-1)/(4t)], \cos^{-1}[(t-1)/(4t)]\}$ , and the value of  $x_+(\mathbf{k})$  at that point is  $(1+3t)^2/(16t)$ .

From the behavior of  $x_{\pm}(\mathbf{k})$  in the neighborhood of the critical points in reciprocal space it is possible to deduce the form of  $G(x)$  near the critical values of  $x$ ,<sup>20</sup> including the value of  $G(0)$  and  $G(1)$ . If the singular part of  $G(x)$  is called  $F(x)$ , then it can be shown that

$$\begin{aligned} F(x) = & B_1 \ln|x-t| + B_2 \ln|x-(1-t)| \\ & + B_3 H[x-(1-t)], \quad 0 < t < \frac{1}{5}, \\ F(x) = & C_1 \ln|x-t| + C_2 \ln|x-(1+3t)^2/16t| \\ & + C_3 H[x-(1-t)], \quad \frac{1}{5} < t < \frac{1}{2}, \end{aligned} \quad (12a)$$

TABLE II. Convergence of the moment-singularity approximating  $G_a^{(N)}(x)$  to the exact distribution of  $G(x)$  with increasing  $N$ . ( $N$  is the number of moments used.)

$x$	$F(x)$	$G_a^{(1)}(x)$	$G_a^{(2)}(x)$	$G_a^{(3)}(x)$	$G_a^{(4)}(x)$	$G_a^{(5)}(x)$	$G_a^{(6)}(x)$	$G(x)$
0	0.450	0.637	0.637	0.637	0.637	0.637	0.637	0.637
0.05	0.528	0.653	0.697	0.689	0.670	0.687	0.689	0.683
0.1	0.620	0.689	0.763	0.752	0.734	0.739	0.739	0.740
0.3	1.36	1.25	1.34	1.34	1.36	1.35	1.35	1.34
1/3								
0.4	1.25	1.08	1.12	1.14	1.16	1.17	1.18	1.17
0.5	1.11	0.897	0.897	0.911	0.911	0.937	0.937	0.951
0.6	1.17	0.939	0.890	0.901	0.879	0.891	0.887	0.908
0.7	1.52	1.28	1.20	1.20	1.17	1.16	1.16	1.14
0.75								
0.9	0.942	0.755	0.682	0.671	0.688	0.694	0.695	0.733
0.95	0.799	0.637	0.599	0.585	0.604	0.621	0.619	0.631
1	0.6851	0.551	0.551	0.551	0.551	0.551	0.551	0.551

<sup>19</sup> We shall assume, as did Montroll in reference 1, that  $t \leq \frac{1}{2}$ .

<sup>20</sup> For a discussion see Sec. III of this paper, also Van Hove, reference 15, Secs. II and III. For comparison see Montroll, reference 1.

where  $H$  is the Heaviside unit function,  $H(y)=0$ , for  $y < 0$ ,  $H(y)=1$ , for  $y > 0$ , and

$$B_1 = -[t(1-2t)]^{-1/2}/\pi^2;$$

$$B_2 = -(1-3t)[t(1-t)(1-2t)(1-5t)]^{-1/2}/\pi^2;$$

$$B_3 = -(2/t\pi^2) \int_0^{\pi/2} d\theta [(1-3t)/t + (1+k^2 \sin^2 \theta)^{1/2}]^{-1},$$

$$k^2 = (3t^2 - 2t + 1)/t^2; \quad (12b)$$

where

$$C_1 = -[t(1-2t)]^{-1/2}/\pi^2;$$

$$C_2 = (16t/\pi^2)(1-t)^{-1}[3(5t-1)(3t+1)]^{-1/2};$$

$$C_3 = [(1-3t)/\pi][t(1-t)(1-2t)(5t-1)]^{-1/2}.$$

The value of  $G(x)$  at 0 and 1 is

$$G(0) = (1+t)[t(1+2t-3t^2)]^{-1/2}/\pi; \quad G(1) = t^{-1/2}/\pi. \quad (12c)$$

In order to apply the moment-singularity method we subtracted  $F(x)$  from  $G(x)$  and approximated the remainder by a linear combination of  $n$  Legendre polynomials. The first  $n-2$  coefficients of the expansion are determined by the requirement that first  $n-2$  moments of the approximate function agree with the exact ones. The remaining two coefficients are then determined by fixing the end points. Hence, when  $N$  moments are known, the approximate distribution function  $G_a^{(N)}(x)$  is

$$G_a^{(N)}(x) = F(x) + \sum_{k=0}^{N+1} A_k^{(N)} P_k(2x-1), \quad 0 \leq x \leq 1, \quad (13a)$$

where

$$A_k^{(N)} = (2k+1) \int_0^1 P_k(2x-1)[G(x)-F(x)]dx \quad (13b)$$

for  $k \leq N-1$ , and

$$\begin{aligned} A_N^{(N)} + A_{N+1}^{(N)} &= G(1) - F(1) - \sum_{i=0}^{N-1} A_i^{(N)}, \\ (-1)^N A_N^{(N)} + (-1)^{N+1} A_{N+1}^{(N)} &= G(0) - F(0) - \sum_{i=0}^{N-1} (-1)^i A_i^{(N)}. \end{aligned} \quad (13c)$$

Note that  $A_k^{(N)}$  is independent of  $N$ , the order of the approximation for  $k \leq (N-1)$ .

When the unmodified method of moments is used to approximate  $G(x)$ , the approximating function, when  $N$  moments are known,  $G_m^{(N)}(x)$  is

$$G_m^{(N)}(x) = \sum_{k=0}^{N-1} B_k^{(N)} P_k(2x-1); \quad (14)$$

the  $B_k^{(N)}$  are again given by Eq. (13b) with  $F(x)$  set equal to zero.

A numerical evaluation of the  $A_k^{(N)}$  and the  $B_k^{(N)}$  was made for  $t=1/3$ . This is the value of  $t$  for which Montroll<sup>1,18</sup> evaluated  $G(x)$  in a closed form, so that

comparisons can be made. This will indicate how useful the moment-singularity method is.

For  $t=1/3$ , Eqs. (12) and (13) yield

$$G_a^{(N)}(x) = -(3/\pi^2) \ln|x - \frac{1}{3}| - (4/\pi^2) \ln|x - \frac{3}{4}| \\ + \sum_{k=0}^{N+1} A_k^{(N)} P_k(2x-1), \quad (15)$$

$$G(0) = 2/\pi; \quad G(1) = \sqrt{3}/\pi.$$

The values of the  $A_k^{(N)}$  in different orders of approximation  $N \leq 6$  are summarized in Table I. The values of the  $B_k^{(N)}$ , which are all independent of  $N$ , are

$$B_0 = 1, \quad B_1 = 0, \quad B_2 = -0.4167, \quad B_3 = 0, \\ B_4 = -0.2269, \quad B_5 = -0.2290. \quad (16)$$

Comparisons between  $G(x)$ ,  $G_a^{(N)}(x)$ , and  $G_m^{(N)}(x)$  are presented in Tables II, III, and IV.

### III. CONCLUSION

As can be seen from Table IV, use of the nonsingular method of moments results in a great increase in the accuracy of the approximation over the usual method of moments for the same number of moments. The additional work required to find  $F(x)$  was small, for  $t=1/3$ . For this particular value of  $t$ ,  $x_{\pm}(\mathbf{k})$  can be expanded in a Taylor series near the critical points, and the behavior of  $G(x)$  near the critical points can be read off directly from Van Hove. For other values of  $t$ , the critical points  $(0,0)$ ,  $(\pi,\pi)$  are of the type which Van Hove calls generalized critical points. The behavior of  $x_{\pm}(k)$  near these points is

$$x(\mathbf{k}) = x_c + |\xi|^2 \psi(\xi/|\xi|) + O(|\xi^3|), \quad \xi = \mathbf{k} - \mathbf{k}_c.$$

To find the form of  $G(x)$  near these points, the integral in Eq. (2) has to be evaluated, terms of  $O(|\xi^3|)$  being neglected. Transforming to polar coordinates in reciprocal space, the integration over the radial variable can be done immediately by means of the delta function. This leaves an integration over the angle variable, whose evaluation may be quite difficult if it cannot be found in the tables. The integral representing  $B_3$  in Eq. (12b) is one such case.

However, even when the exact form of the singular part of  $G(x)$ , such as the coefficients multiplying the logarithmic terms in two dimensions, or the square root terms in three dimensions, are not known, convergence would be improved greatly if the approximating function contained terms of the correct singular form. The coefficients of these terms could then be determined by the moments. Thus, for the case considered in this paper, it would have been preferable if the approximating function  $G_a(x)$  consisted of polynomials multiplying the logarithmic terms whose value at the critical points is the correct one. This would have taken account of terms in the distribution func-

TABLE III. Convergence of the unmodified moment approximation  $G_m^{(N)}(x)$ , to the exact distribution of  $G(x)$  with increasing  $N$ . ( $N$  is the number of moments used.)

$x$	$G_m^{(1)}(x)$	$G_m^{(2)}(x)$ $=G_m^{(3)}(x)$	$G_m^{(4)}(x)$ $=G_m^{(5)}(x)$	$G_m^{(6)}(x)$	$G(x)$
0	1	0.583	0.332	0.103	0.637
0.05	1	0.702	0.655	0.646	0.683
0.1	1	0.808	0.861	0.770	0.740
0.3	1	1.11	1.13	1.20	1.34
1/3	1	1.14	1.14	1.21	$\infty$
0.4	1	1.18	1.13	1.20	1.17
0.5	1	1.21	1.12	1.12	0.951
0.6	1	1.18	1.13	1.06	0.908
0.7	1	1.11	1.13	1.07	1.14
0.75	1	1.05	1.12	1.09	$\infty$
0.9	1	0.808	0.861	0.953	0.733
0.95	1	0.702	0.655	0.664	0.631
1	1	0.583	0.332	0.561	0.551

tion of the form  $(x-x_c) \ln|x-x_c|$ . A smaller number of moments might then have been used to obtain the same accuracy. This was not done because of the extra work involved in solving simultaneous equations for the coefficients of the multiplying polynomials since no orthogonal set would be available. In general, the amount of work required to find more moments has to be balanced against the work involved in solving simultaneous equations.

There is a theorem of Feynman<sup>21</sup> which is useful in finding *some* critical points without diagonalizing  $\mathbf{M}(\mathbf{k})$ . This theorem states that, if  $\mathbf{A}(\alpha)\Psi(\alpha) = \lambda(\alpha)\Psi(\alpha)$  and  $\mathbf{A}(\alpha)$  is Hermitian, then

$$\frac{\partial \lambda(\alpha)}{\partial \alpha} = \left( \Psi(\alpha), \frac{\partial \mathbf{A}(\alpha)}{\partial \alpha} \Psi(\alpha) \right).$$

Thus  $|\text{grad} \omega^2(k)| = 0$  whenever  $|\text{grad} \mathbf{M}(\mathbf{k})| = 0$ . This can happen only at those  $\mathbf{k}$  where all the branches have critical points simultaneously.

A great difficulty, in many cases, in finding the analytic form of the distribution, is the location of *all* the critical points. Van Hove's arguments predict only

TABLE IV. Comparison of the unmodified approximation  $G_m^{(6)}(x)$ , the moment singularity approximation  $G_a^{(6)}(x)$ , and the exact distribution  $G(x)$  when six moments are used.

$x$	$G_m^{(6)}(x)$	$G_a^{(6)}(x)$	$G(x)$
0	0.103	0.637	0.637
0.05	0.646	0.689	0.683
0.1	0.770	0.739	0.740
0.3	1.20	1.35	1.34
1/3	1.21	$\infty$	$\infty$
0.4	1.20	1.18	1.17
0.5	1.12	0.937	0.951
0.6	1.06	0.887	0.908
0.7	1.07	1.16	1.14
0.75	1.09	$\infty$	$\infty$
0.9	0.953	0.695	0.733
0.95	0.664	0.619	0.631
1	0.561	0.551	0.551

<sup>21</sup> R. P. Feynman, Phys. Rev. **56**, 340 (1939).

the minimum number of critical points but give no upperbound. It is generally easy to find those which are at symmetry points of the Brillouin zone. In most cases investigated thus far,<sup>1-3</sup> these are the only critical points. This might be due to the assumption made in these cases that the forces are of short range. The number of critical points might be expected to increase with the range of the interaction, and their location is then more likely to be at nonsymmetry points of the zone. Rosenstock<sup>17</sup> has devised a method for examining the presence of critical points inside the zone from the behavior of  $\omega^2(\mathbf{k})$  on the boundary.

For three dimensions, with short-range forces, when the critical points are at the symmetry points the matrix  $\mathbf{M}(\mathbf{k})$ , is easy to diagonalize at these points to find the critical frequencies. To find the exact form of the distribution near the critical points  $x_c$ , degenerate perturbation theory has to be used to find the eigenvalues in the vicinity of  $\mathbf{k}_c$ . However, since in three dimensions  $G(x)$  does not become infinite at  $x_c$ , it might be sufficient to use the right form of  $G(x)$  near  $x_c$ ; the coefficients of the singular part would then be determined by the moment as mentioned above. This would agree with the results of Rosenstock.<sup>17</sup>

## Electrical Properties and the Solid-Vapor Equilibrium of Lead Sulfide

ROBERT F. BREBRICK AND WAYNE W. SCANLON  
U. S. Naval Ordnance Laboratory, White Oak, Maryland  
(Received June 21, 1954)

Natural PbS crystals were exposed to various pressures of sulfur-vapor near 500°C and then quenched. Calculations based on the penetration of *p-n* junctions gave an interdiffusion constant of  $2 \times 10^{-6}$  cm<sup>2</sup>/sec at 550°C. The temperature dependence of the Hall coefficient and resistivity for several treated crystals was determined. A forbidden energy gap of  $0.37 \pm 0.01$  ev and an electron to hole mobility ratio of 1.4 was obtained.

### I. INTRODUCTION

AT low temperatures the electrical properties of ionic or partially ionic binary compound semiconductors may be attributed to the presence of donor and acceptor levels produced either by foreign atoms or by deviations from stoichiometric proportions. In the absence of appreciable concentrations of foreign atoms the electrical properties are dependent to a large extent upon deviations from stoichiometry. Current statistical models predict that it is possible for all crystalline ionic compounds to show these deviations from stoichiometry.<sup>1</sup> The crystal can exist as a single phase over a range of composition through the inclusion of interstitial atoms and/or vacant lattice sites. While this range of composition is generally too small to be detected chemi-

cally it is revealed through variations in electrical properties which are composition sensitive.

The ability of the crystal to exist as one phase over a range of composition implies some thermodynamic consequences which are important in the preparation of these materials with desired electrical properties. In the case of lead sulfide the equilibrium system, crystal-vapor, has two phases and two components, lead and sulfur. According to the Gibbs phase rule this system possesses two degrees of freedom. Hence when the temperature and vapor pressure of either component are fixed, all of the intensive properties of the system, such as the composition of the crystal and vapor phases, are fixed also. The vapor phase will in general contain two components in a different proportion than the crystal.<sup>2</sup> Therefore, under equilibrium conditions, the crystal can be held at a fixed composition over a range of temperature only by appropriately adjusting the vapor pressure of one of the components or the total pressure. On the other hand, if one of these pressures is held constant and the temperature varied or *vice versa*, then the composition of the crystal will change in general.

In principle, therefore, the composition of a crystalline binary compound can be varied by heat treatments in a controlled atmosphere comprised of its components. In practice, it is necessary to work at those tempera-




FIG. 1. Apparatus used for treating crystals.

<sup>1</sup> R. H. Fowler and E. A. Guggenheim, *Statistical Thermodynamics* (Cambridge University Press, Cambridge, 1949), Chap. XII.

<sup>2</sup> C. Wagner, *Thermodynamics of Alloys* (Addison-Wesley Press, Inc., Cambridge, 1952), p. 67.