

Polymorphic Transition in LiH†

DAVID P. SCHUMACHER

Department of Chemistry, Tufts University, Medford, Massachusetts

(Received January 29, 1962)

The transition pressure at which LiH would be expected to go from a NaCl to a CsCl structure is calculated to be 3050 and 3800 kg/cm² by using two different potential functions. Also the transition pressures are calculated for KCl and RbI to be 22 500 and 5400 kg/cm², respectively, which are in good agreement with the observed values.

A QUANTUM-MECHANICAL calculation of the metallic transition pressure in LiH assuming no initial transition to the denser CsCl structure has been carried out by Behringer.¹ It is possible that this assumption is unwarranted.

The variation of the molecular volume with pressure is given by

$$\dot{p} = -\frac{\partial E_{\text{mol}}}{\partial v} = -\frac{\partial E_{\text{mol}}}{\partial a} \frac{da}{dv},$$

where a is the nearest Li-H distance and $E_{\text{mol}} = L\varphi$, L being Avogadro's number. The potential function² φ is taken as

$$\varphi = A \exp(-a/\rho) - \alpha e^2/a, \quad (1)$$

or as

$$\varphi = A/a^n - \alpha e^2/a, \quad (2)$$

where α is the Madelung constant, ρ is a constant which for saline metal hydrides is equal to 0.465×10^{-8} cm,^{3,4} n is the Born exponent taken as 5.0 for LiH,⁵ and A is found from $(\partial \varphi / \partial a)_{a=a_0} = 0$. These equations then lead to

$$\dot{p} = \frac{\alpha e^2}{3Ka^2} \left[\frac{e^{(a_0-a)/\rho}}{a_0^2} - \frac{1}{a^2} \right] \quad (3)$$

and

$$\dot{p} = \frac{\alpha e^2}{3Ka^3} \left[\frac{a_0^{n-1}}{a^n} - \frac{1}{a} \right], \quad (4)$$

from Eqs. (1) and (2), respectively. The term K is a structure factor.⁶ The values of a_0 for LiH are 2.04 Å for

TABLE I. Summary of results.

	Obs. ^a	Transition pressures (kg/cm ²)			$\Delta v/v_0$	
		Calc.	Calc. by others		Obs. ^a	Calc.
KCl	20 060	22 500 ^b	74 000 ^d	31 400 ^e	0.085	0.080
RbI	4050	5400 ^b	22 000 ^d		0.035	0.032
LiH	...	3050 ^b	0.0035
		3800 ^c				0.0039

^a P. W. Bridgman, Proc. Am. Acad. Arts Sci. **76**, 1 (1948).

^b From Eq. (3).

^c From Eq. (4).

^d R. B. Jacobs, Phys. Rev. **54**, 468 (1938).

^e See reference 6.

the NaCl structure⁷ and 2.12 Å for a CsCl structure. The a_0 for the CsCl case is calculated from the sum of the tabular radii⁸ with a coordination number correction for the cation.⁹

Similar calculations were also extended to KCl and RbI using the appropriate value of $\rho = 0.345$ Å. The results are tabulated in Table I.

The value of ρ used is that which gives the best agreement between calculated and observed lattice energies⁵ and cannot be expected to express the repulsive energy adequately at high pressures. The agreement between calculated and observed transition pressures is quite good considering the empirical nature of ρ .

If ρ is increased from 0.345 to 0.374 Å in the case of RbI, the transition pressure will be increased to 5860 kg/cm², while for LiH, if $\rho = 0.345$ Å, the transition pressure will be decreased to 2400 kg/cm². For an uncertainty of ± 0.01 Å in ρ , the transition pressure will vary by ± 50 for LiH and ± 230 kg/cm² for RbI.

† Work supported by the U. S. Atomic Energy Commission.

¹ R. E. Behringer, Phys. Rev. **113**, 787 (1959).

² M. Born and J. E. Mayer, Z. Physik **75**, 1 (1932).

³ T. R. P. Gibb, Jr., J. Inorg. & Nuclear Chem. (to be published).

⁴ E. C. Baughan, Trans. Faraday Soc. **55**, 736 (1959).

⁵ J. Sherman, Chem. Revs. **11**, 93 (1932).

⁶ Per-Olov Löwdin, *Advances in Physics*, edited by N. F. Mott (Taylor and Francis, Ltd., London, 1956), Vol. 5, p. 1.

⁷ E. Zintl and A. Harder, Z. physik. Chem. **B14**, 265 (1931).

⁸ T. R. P. Gibb, Jr. and D. P. Schumacher, J. Phys. Chem. **64**, 1407 (1960).

⁹ C. Kittel, *Introduction to Solid-State Physics* (John Wiley & Sons, Inc., New York, 1959), p. 81.