

should read

$$g_0 \epsilon Z^{-1} [\mathbf{m}_V m_N]^{1/2} [2\Omega]^{-1/2} [\omega_N(\mathbf{p}-\mathbf{k}) \omega_\theta(\mathbf{k}) \omega_V(\mathbf{p})]^{-1/2} \\ \times u_N(\mathbf{p}-\mathbf{k}) \{i(\mathbf{p}-\mathbf{k}) + m_N\}^{-1} \beta Q_0 u_V(\mathbf{p}).$$

**Momentum Distributions for Protons in  $\text{Li}^6$  in the Cluster Model**, J. STRNAD [Phys. Rev. **125**, 1639 (1962)]. Owing to a calculational mistake the main conclusion of the paper, i.e., that the absence of a dip in the momentum distribution for the least bound protons in  $\text{Li}^6$  at zero momentum transfer can be explained in the cluster model, is invalid. In fact, the cluster model as well as the shell model does give the dip for zero momentum transfer for the least bound protons. Thanks are due to Dr. G. Jacob and Dr. P. Said for pointing this out to the author.

**3d Band Structure of Cr**, M. ASDENTE AND J. FRIEDEL [Phys. Rev. **124**, 384 (1961)]. In this paper, energies in the theoretical calculations are given in Rydberg instead of in atomic units as would have been correct.

Consequently, units should be changed in Figs. 2, 3, 4, 5, 7. Only Fig. 7(b) needs to be modified




FIG. 7(b). Density of states curves  $g(E)$ . Open circles:  $g(E_F)$  deduced from experimental values of  $C_v$ ; solid points:  $g(E_F)$  deduced from experimental values of  $\chi$ .

and is given here again; the general conclusions are unaffected, however.

**Neutron Coherent Scattering Amplitudes for Cd and Eu**, G. ARNOLD AND N. NERESON [Phys. Rev. **124**, 1848 (1961)]. Errors have been discovered in

the measured transmissions of the  $\text{Eu}_2\text{O}_3$  and  $\text{CdO}$  samples which were used to determine the neutron coherent scattering amplitudes of Eu and Cd. A first indication that the reported values of  $f_{\text{Eu}}$  might be in error was a confirmative experiment using  $\text{EuO}$  which yielded a value for  $f_{\text{Eu}}$  in disagreement with that obtained from  $\text{Eu}_2\text{O}_3$ . Subsequently, repeated transmission measurements made in the direct beam from the Pb monochromating crystal were found to be in error for some of the oxide samples. These transmission measurements are now made in the scattered beam from the nickel standard set to a diffraction peak. This method reduces background effects associated with an impure neutron beam and gives measured transmissions in good agreement with calculated values.

The corrected values of  $F^2$  in the two separate measurements on  $\text{CdO}$  are shown in Fig. 1. The




FIG. 1. Graph of square of intensity structure factors vs  $(\sin\theta/\lambda)^2$  for  $\text{CdO}$ .

averaged extrapolated value of  $F^2$  at  $0^\circ$  scattering angle is  $(19 \pm 1.5) \times 10^{-24} \text{ cm}^2$ . Using  $0.58 \times 10^{-12} \text{ cm}$  as the scattering amplitude for oxygen, one obtains a value of  $f_0(\text{Cd}) = (0.71 \pm 0.04) \times 10^{-12} \text{ cm}$  for the nonresonant coherent scattering length of Cd; the absolute value of the scattering amplitude at  $\lambda = 1.391 \text{ \AA}$  is  $|f_{\text{Cd}}| = (0.51 \pm 0.03) \times 10^{-12} \text{ cm}$ . This value is still in some disagreement with that obtained by Peterson and Smith<sup>1</sup> from their single-crystal work on  $\text{CdI}_2$  which gives a value of  $|f_{\text{Cd}}| = 0.42$  at  $\lambda = 1.39 \text{ \AA}$ .<sup>2</sup>

Corrected scattering amplitudes from the  $\text{Eu}_2\text{O}_3$  measurements are  $f_0(\text{Eu}) = (0.85 \pm 0.05) \times 10^{-12} \text{ cm}$  and  $|f_{\text{Eu}}| = (0.60 \pm 0.04) \times 10^{-12} \text{ cm}$  at  $\lambda = 1.391 \text{ \AA}$ . Recent measurements from the compound  $\text{EuO}$  give  $f_0(\text{Eu}) = (0.88 \pm 0.04) \times 10^{-12} \text{ cm}$  and  $|f_{\text{Eu}}| = (0.63 \pm 0.03) \times 10^{-12} \text{ cm}$ .

<sup>1</sup> S. W. Peterson and H. G. Smith, Phys. Rev. Letters **6**, 7 (1961).

<sup>2</sup> S. W. Peterson (private communication).