

fitted to a theoretical relation.¹⁶ It contains an outer Coulomb correction using $r_0=\lambda_c$. This value for α_3/η and the present experimental a give $\alpha_1/\eta=0.184$. The same α_3/η and a , except using $r_0=0.5\lambda_c$, give $\alpha_1/\eta=0.174$. If we now apply the inner Coulomb correction as well, we get $\alpha_1/\eta=0.170$ and 0.172 using $r_0=\lambda_c$ and $0.5\lambda_c$, respectively.

The Hamilton and Woolcock analysis gives $\alpha_1/\eta=0.178\pm0.005$ with outer Coulomb correction using $r_0=\lambda_c$. The comparable value found from this experiment is $\alpha_1/\eta=0.184$ using the α_3/η of Hamilton and Woolcock. The present experiment is therefore not in

¹⁶ The value of α_3/η preferred by Hamilton and Woolcock is not directly comparable with that reported by Fischer and Jenkins. When the Fischer and Jenkins result is corrected using $r_0=\lambda_c$, it becomes $\alpha_3/\eta=-0.108$. Furthermore, the Fischer and Jenkins amplitude fell off at low energies by $1\frac{1}{2}$ standard deviations.

disagreement with the consistent low-energy pion parameters of Hamilton and Woolcock.

V. ACKNOWLEDGMENTS

The author wishes to thank Professor Jack Steinberger for suggesting this experiment and for continued guidance. The entire Columbia Bubble Chamber group carried out the exposure. The conscientious work of many scanners is appreciated, notably Irene Garder, Alex Rytov, and Mary Johnson, and the careful measurements and analysis by Herman Kaufman. The author also thanks Dr. Daniel Tycko for assistance in the use of the IBM 650.

Special appreciation is expressed to Dr. Edgar Jenkins and Dr. Gerhard Fischer for discussions of their similar experiment at the outset of this experiment

Probable Example of the Decay $\Sigma^0 \rightarrow \Lambda + e^+ + e^-$ in Nuclear Emulsion

D. H. DAVIS,* R. LEVI SETTI, M. RAYMUND, AND G. TOMASINI†

The Enrico Fermi Institute for Nuclear Studies,‡ The University of Chicago, Chicago, Illinois

(Received March 15, 1962)

An event has been observed in nuclear emulsion which is attributed to the production of a Σ^0 hyperon and its subsequent decays via the mode $\Sigma^0 \rightarrow \Lambda + e^+ + e^-$. From the apparent concurrence of the tracks involved in the disintegrations, it has been deduced that the Σ^0 lifetime for this event is $<10^{-14}$ sec.

THE lifetime of the Σ^0 hyperon is expected to be within the range 10^{-18} – 10^{-20} sec,^{1,2} and possible methods for its determination have been recently discussed by Dreitlein and Primakoff.³ Until now the sole experimental value reported⁴ was obtained from the observation of Σ^0 hyperon decay events in a hydrogen bubble chamber by Alvarez *et al.*⁵ The Σ^0 hyperon lifetime was quoted to be less than 10^{-11} sec. During the systematic scanning for K^- meson captures at rest in an Ilford K5 emulsion stack an event has been found which is attributed to the decay of a Σ^0 hyperon via the Dalitz mode, i.e.,

$$\Sigma^0 \rightarrow \Lambda^0 + e^+ + e^- \quad (1)$$

* Fulbright Travel Scholar on leave from University College London.

† On leave of absence from the University of Genoa.

‡ Work supported by the U. S. Air Force Office of Scientific Research, Contract AF 49(638)-209.

¹ M. Gell-Mann and A. H. Rosenfeld, *Ann. Rev. Nuclear Sci.* **7**, 407 (1957).

² J. Dreitlein and B. Lee, *Phys. Rev.* **124**, 1274 (1961).

³ J. Dreitlein and H. Primakoff (to be published).

⁴ W. H. Barkas and A. H. Rosenfeld, University of California Radiation Laboratory Report UCRL-8030, 1961 (unpublished), revised.

⁵ L. W. Alvarez, H. Bradner, P. Falk-Vairant, J. D. Gow, A. H. Rosenfeld, F. T. Solmitz, and R. D. Tripp, University of California Radiation Laboratory Report UCRL-3775, 1957 (unpublished).

This decay mode was first observed by Eisler *et al.*⁶ in a hydrogen bubble chamber. The high spatial resolution of the emulsion technique has enabled a much more precise experimental upper limit to be placed upon the decay length and hence the lifetime of the Σ^0 hyperon.

A photograph of the event to be described is shown in Fig. 1 and the measurements made on it are summarized in Table I. A K^- meson is captured at point O and three fast charged particles are seen to be emitted from the resulting nuclear disintegration. One is due to a charged π meson, probably negative, because it interacts in flight after a path length of 9.1 mm to form a star from which no charged secondaries are emitted. The energy of this π meson, as determined by ionization and scatter-

TABLE I. Summary of the measurements.

Track	φ (deg)	λ (deg)	Observed length (μ)	g^*	$p\beta$ MeV/c	Kinetic energy (MeV)
e_1	184.8	+10.6	2200	$0.96 \pm .04$	28 ± 9	28 ± 9
e_2	187.7	+ 5.7	3300	$1.07 \pm .03$	48 ± 10	48 ± 10
π^\pm	45.3	+16.2	9100	$1.28 \pm .02$	90 ± 17	57 ± 4

⁶ F. Eisler, R. Plano, N. Samios, J. Steinberger, and M. Schwartz, *Phys. Rev.* **110**, 226 (1958).





FIG. 1. Photomicrograph of the event.

ing measurements along its track, is (57 ± 4) MeV. Tracks e_1 and e_2 are due to an electron pair of opening angle 5.1° . The energies of the electrons, as measured from multiple Coulomb scattering, are (48 ± 10) MeV and (28 ± 9) MeV, respectively. The energy of the pair is thus (76 ± 14) MeV. At the vertex of the capture star there is also a small blob, which is probably either the track of a heavy recoiling nucleus or of a slow electron. A mass measurement of the primary particle, which entered the stack in the direction of the beam, was made using the constant sagitta scattering method. The value obtained, $(0.8 \pm 0.2) m_K$, clearly indicates that the event is indeed due to a K^- meson capture.

By far the most likely interpretation of this event is that the K^- meson is captured on a bound neutron to form a Σ^0 hyperon



and that the Σ^0 hyperon subsequently decayed via the Dalitz mode. This mode is expected to occur with a frequency of once for every 160–180 normal radiative

Σ^0 hyperon decays^{7,8}. It can be shown that alternative interpretations of this event, viz., the decay of a K^- meson almost at rest, via the mode $K^- \rightarrow \pi^- + \pi^0 + \pi^0$, and the π^0 decay of an unobserved hyperfragment, are extremely improbable.

The upper limit of the decay length of the Σ^0 hyperon was measured in a way similar to that used by other emulsion workers^{9–11} when estimating the π^0 meson lifetime. The direction of the resultant of the electron pair momenta was found to be concurrent with the intersection of the π meson and K^- meson tracks, to within $(0.1 \pm 0.3) \mu$ in the emulsion plane, and to within $(0.4 \pm 0.5) \mu$ in the vertical plane. In this emulsion, Ilford K5, the mean diameter of the developed grains is $(0.55 \pm 0.03) \mu$, whereas in Ilford L4 fine grain emulsion (used for the π^0 meson lifetime estimates by Glasser *et al.*¹⁰) the mean diameter of the developed grains is about 0.35μ . Furthermore, whilst the velocity of the π^0 meson from $K_{\pi^2}^+$ meson decays at rest is precisely known, that of the Σ^0 hyperon in this case is much lower and less well determined. Both of these factors render one unable to measure the decay length of the Σ^0 hyperon with the same precision as can be achieved for individual π^0 mesons. However, Gilbert *et al.*¹² and the European K^- Collaboration¹³ have shown that for K^- -meson captures on bound protons in which a Σ hyperon and a π meson are the sole charged particles emitted, the sum of their kinetic energies is usually some 15 MeV less than the Q value for the reaction on a free proton. For a π meson energy of about 57 MeV, the most likely value for the Σ^0 hyperon energy is about 30 MeV, corresponding to a $\beta = 0.22$. The upper limit of the decay time of the Σ^0 hyperon is then $\sim 4 \times 10^{-15}$ sec. Even if the energy of the Σ^0 hyperon is assumed to be as low as 6 MeV, β_{Σ^0} would become 0.1 and the upper limit of the decay time can still be set at $\sim 1 \times 10^{-14}$ sec.

We are indebted to Professor V. L. Telegdi for helpful discussions and we wish to express our gratitude to Dr. E. J. Lofgren and the Bevatron staff for the exposure to the separated K^- beam.

⁷ G. Feinberg, Phys. Rev. **109**, 1019 (1958).

⁸ G. Feldman and T. Fulton, Nuclear Phys. **8**, 106 (1958).

⁹ G. Harris, J. Orear and S. Taylor, Phys. Rev. **106**, 327 (1957).

¹⁰ R. G. Glasser, N. Seeman, and B. Stiller, Phys. Rev. **123**, 1014 (1961).

¹¹ R. F. Blackie, A. Engler, and J. H. Mulvey, Phys. Rev. Letters **5**, 384 (1960).

¹² F. C. Gilbert, C. E. Violet, and R. S. White, Phys. Rev. **107**, 228 (1957).

¹³ K^- European Collaboration Part I, Nuovo cimento **13**, 690 (1959).




FIG. 1. Photomicrograph of the event.